

Production of Protease by *Bacillus* sp. N-40 Isolated from Soil and Its Enzymatic Properties

Nihan Sevinc and Elif Demirkan*

¹Uludag University, Faculty of Arts and Sciences, Department of Biology, Bursa, TURKEY

ABSTRACT

Fifty four *Bacillus* strains isolated from different soils were screened for protease production. One strain producing maximum protease was selected. In order to enhance the production of protease, the effects of major medium ingredients, such as carbon, nitrogen sources and metal ions, on the production of the enzyme were investigated. Among the carbon sources used, fructose showed the highest potential for the production. The best organic nitrogen source was skim milk. Inorganic nitrogen sources were not as effective as organic sources. Addition of combine metal ions minimized the enzyme production. Combinations of Ca²⁺ and Mg²⁺ in medium were the best. Both ions were not effective alone. Increased production (51%) of the enzyme was obtained by manipulating the medium composition. The optimum pH and temperature for the purified enzyme activity were 7.0 and 55°C, respectively. The study of its stability showed that the enzyme is stable in the alkaline pH range 6.0-9.0, and at temperatures between 40 and 70 °C. The enzyme was also thermostable (77% at 55 °C for 3 h). The enzyme activity was stimulated by Mn²⁺ and Ca²⁺. It showed a single band with a molecular weight of 52 kDA by SDS-PAGE.

Key words: *Bacillus*, Isolation, Protease, Partial purification, Characterization

INTRODUCTION

Proteases refers to a group of enzymes whose catalytic function is to hydrolyze proteins. They are also called proteolytic enzymes or proteinases. Proteases are classified according to their structure or the properties of the active site. There are several kinds of proteases such as serine-, metallo-, carboxyl-, acidic-, neutral-, and alkaline proteases. Proteases are one of the most important group of industrial enzymes, and commercial proteases account for nearly 60% of the total industrial enzyme market. They are widely used in leather processing, detergent industry, food industries, bioremediation process, pharmaceutical, textile industry, waste processing companies, and in the film industry etc. (Rao et al. 1998). Proteases are obtained from plants and animal organs and microorganisms, with the majority obtained from bacteria and fungi. Currently, a large proportion of commercially available proteases are derived from *Bacillus* strains (Mehratto et al. 1999). *Bacillus* proteases are predominantly extracellular and can be concentrated in the fermentation medium. The most studies focused on screening proteases with a criterion set only to increase activity levels (Kim et al. 1994). Selection of the right organism plays a key role in high yield of desirable enzymes. On the other hand, it is a well-known fact that extracellular protease production in microorganisms is greatly influenced by media components, especially carbon and nitrogen sources (McKeller and Cholette 1984), metal ions (Adinarayana et al. 2003). Isolation and characterization of new promising strains using carbon and nitrogen source is a continuous process (Asokan and Jayanthi 2010).

In the present study, *Bacillus* sp. strains isolated from 31 different Turkish soils were screened for proteolytic activity. A strain which have the highest proteolytic activities were determined and some fermentation conditions such as carbon and nitrogen sources and various metal ions on protease production were investigated. A modified medium was defined for high production of protease. Protease was partially purified and characterized.

MATERIALS AND METHODS

Bacterial isolation and screening for proteolytic activity

Soil samples of 31 different cities of Turkey were collected, serially diluted and plated on nutrient agar plates. The isolated pure colonies were examined for *Bacillus* with standart description of Bergey's Manual of Determinative Bacteriology (Buchanan and Gibbons 1974). A total of 54 *Bacillus* sp. were screened for protease production by using casein agar (0.5% casein in nutrient agar) and skim milk agar (1% skim milk in nutrient agar). Proteolytic activities of *Bacillus* sp. were detected on the basis of appearance of clear zones around the bacterial colonies. The protease positive colonies were obtained a pure cultures and the most potent four strains were selected for quantitative test of protease.

* Corresponding author: edemirkan@uludag.edu.tr

Protease production

Two culture media used for protease production and compared. One of them contained (% w/v) : glucose-0.1, peptone-1, yeast extract-0.02, MgSO₄-0.01, CaCl₂-0.01, K₂HPO₄-0.05 (pH 7.0) (Qadar et al. 2009). Other composed of (% w/v) : glucose-1, peptone-0.5, yeast extract-0.3, MgCl₂-0.02, CaCl₂-0.04 (pH 7.0) (Sangeetha et al. 2008). Glucose was sterilized separately and aseptically added to the flasks containing the liquid medium, after cooling. The precultures were cultivated in Nutrient Broth medium (0.8% w/v) for 18 h. Then, overnight cultures with OD₆₀₀=0.3 were inoculated at 1% in enzyme production media (150 mL in 500 mL Erlenmeyer flasks) and incubated at 37 °C for 16, 24, 40, 48, 64 and 72 h in a shaking incubator (150 rpm). At the end of each period, the cultures were centrifuged (6000 rpm, 10 min) and the supernatants were used for determination of proteolytic activity. Bacterial biomass was determined by measuring optical density at 600 nm.

Capabilities of protease production of four strains were examined, and one strain and the best medium were selected for further studies.

Assay of proteolytic activity

Total protease activity was measured using a casein substrate by a modification of the Anson Method (Keay and Wildi 1970). A 1 ml of the culture supernatant was mixed with 1 ml 0.05 M phosphate buffer-0.1 M NaOH (pH 7.0 adjusted with phosphoric acid) containing 2% casein, and incubated for 10 min at 37 °C. The reaction was stopped by adding 2 ml 0.4 M Trichloroacetic acid. After 30 min stand at room temperature, the precipitate was removed by centrifugation and the optical density of the assays was measured at 660 nm. A standard curve was generated using solutions of 0–60 µg/mL tyrosine. One unit of protease activity was defined as the amount of enzyme required to liberate 1 µg/mL tyrosine under the experimental conditions used.

Effect of nutrients on enzyme production

The effects of some nutrients such as carbon, nitrogen sources and some metal salts on enzyme production were investigated. Glucose (0.1% w/v) was replaced in the production medium with maltose, sucrose, fructose, glycerol and starch. Different organic and inorganic nitrogen sources including corn steep liquor, skim milk, triptone, soybean, casein, (NH₄)₂SO₄ and KNO₃ were tested. These nitrogen sources were used to replace peptone and yeast extract which were the original nitrogen source in growth medium. The culture medium was supplemented with the following metal salts (0.01% w/v): CaCl₂ with LiSO₄, BaCl₂, MnCl₂, CuSO₄; MgSO₄ with LiSO₄, BaCl₂, MnCl₂, CuSO₄. Incubation periods were set as 16, 24, 40, 48, 64 and 72 h.

A new medium including the best source of carbon, nitrogen and metal salts for protease production were improved, and bacteria were grown in this new medium.

Partial purification of crude protease and characterization

The crude protease extract was partially purified by ammonium sulphate (75%) precipitation followed by dialysis. The precipitate was collected by centrifugation at 12,500 rpm for 20 min at 4 °C, and dissolved in 0.01 M phosphate buffer (pH 7.0). The solution was dialyzed against the same buffer at 4 °C for 8 h with 3 changes of the dialysis buffer, and protein concentration was measured as the method of Lowry et al. (1951) using BSA as standard protein.

After partial purification, the enzyme was characterized with its optimal values of pH (4.0-9.0), temperature (37-80 °C), and their stabilities were presented. The effects of various metal ions (Ca²⁺, Mg²⁺, Mn²⁺, Li²⁺, Ba²⁺, Cu²⁺, Zn²⁺) on enzyme activity were determined, and enzyme was incubated with metal ions of 5 mM. Relative activities were expressed as a percentage of the activity of the untreated control taken as 100%.

The approximate molecular weight of partially purified enzyme was determined by Sodium Dodecyl Sulphate Polyacrylamide Gel Electrophoresis (SDS-PAGE) (Laemmli 1970).

RESULTS AND DISCUSSION

Screening and isolation of proteolytic bacteria

Fifty four (54) *Bacillus* sp. strains isolated from soil samples were determined on the basis of morphological and biochemical characteristics. The proteolytic activities of all strains were assayed using skim milk agar and casein agar, and exhibited as diameter of clear zone in mm. Skim milk agar was the best than casein agar

for Qualitative test of protease. Among *Bacillus* sp. strains, six isolates showed high proteolytic activity (18-14 mm), twenty seven isolates showed moderate activity (10-12 mm), fifty five isolates showed weak activity (8-6 mm), and six isolates exhibited very weak activity (4 mm). Four high proteolytic strains (N-3, N-10, N-18, N-40) were checked for Quantitative test of protease in two liquid media. Among tested media, medium of Quadar et al. (2009) was the best than other. In this medium, the maximum protease activity (224 U/ml) was attained after 64 h by *Bacillus* sp. N-40 strain (Table 1). It found that maximum production occurred at end of exponential phase.

Table 1. Effect of incubation time on protease production by four *Bacillus* strains during cell growth in medium of Quadar et al. (2009). Values are shown as means of triplicates.

Incubation time (h)	N-3		N-10		N-18		N-40	
	U/ml	OD ₆₀₀	U/ml	OD ₆₀₀	U/ml	OD ₆₀₀	U/ml	OD ₆₀₀
16	50	0.6	24	0.3	52	0.7	83	0.6
24	64	0.6	56	0.5	68	0.9	100	0.7
40	90	0.8	70	0.9	74	1.0	150	0.8
48	99	0.9	85	1.3	90	1.2	182	0.9
64	70	1.0	80	1.1	98	1.1	224	1.1
72	64	1.1	60	0.8	83	0.7	214	1.0

Other three strains showed low production of protease, and *Bacillus* sp. N-40 which is high capability was selected for further studies.

Effect of carbon, nitrogen and metal ions on protease production

Enzyme formation is largely dependent on the condition of growth of the culture and composition of nutrient medium. The present investigation was aimed at optimization of medium components which have been predicted to play a significant role in enhancing the production of proteases (Gupta et al. 2002).

The research was focused on the improvement of protease level. Various carbon, nitrogen sources and metal ions were used for the production of protease by *Bacillus* sp. N-40. When glucose in basal medium was replaced by five various sugars (maltose, sucrose, fructose, glycerol, starch), fructose was the best source for protease production (Fig. 1). Fructose (278 U/ml) increased the production of protease by 24% when compared to control (glucose, 224 U/ml). The protease production was affected by carbon sources in the order: fructose > sucrose > maltose > control > starch > glycerol. The increase in enzyme production was parallel to the growth rate (64 h). Glycerol caused reduced growth and protease synthesis.




Figure 1. Protease production by *Bacillus* sp. N-40 in various carbon and nitrogen sources.

Reserachers have been reported different results for the best carbon sources. In a similar study, it showed that fructose was the best carbon source for protease production (Mabrouk et al. 1999; Sangeetha et al. 2008).

On the other hand, Starch, sucrose, and lactose proved appreciably good for the protease production by *Bacillus* sp. K-30 (Naidu and Devi 2005). Alkaline protease production was best in wheat bran, starch, glucose and dextrin by *Bacillus licheniformis* and *Bacillus coagulans* (Asokan and Jayanthi 2010). Lactose was the best carbon source that induced the production of protease by *Bacillus subtilis* (El-Safey and Abdul-Raouf 2004). Madzak et al. (2000) recorded that the sucrose is good substrate for production extracellular proteases. Glucose was found to be the optimum carbon source for protease activity by all the four *Bacillus* isolates followed by sucrose, fructose, maltose, starch and cellulose (Boominadhan et al. 2009). However, other works reported that glucose drastically inhibited protease production (Fukushima et al. 1989; Puri et al. 2002). On the other hand, in this study, starch gave lower growth but higher protease production, but Hadj-Ali et al. (2007) has been reported opposite result. Maltose, starch and cellobiose caused low protease production (Shafee et al. 2005), while starch caused high level of enzyme expression in *Bacillus* species (Johnvesly and Naik 2001).

Nitrogen sources also affected enzyme production. Effects of various organic and inorganic nitrogen sources on production of protease were investigated. Results obtained showed that the best nitrogen source for protease production was skim milk (320 U/ml) at 64 h, and enzyme yield was 43% compare to control (224 U/ml). Enzyme production in the presence of casein (250 U/ml) reached at 24 h. Maximum enzyme production were skim milk > casein > control > tryptone > soy bean > corn steep liquor (Fig 1). The increase in enzyme production was parallel to the growth rate for all nitrogen sources. Inorganic sources were found to have no effect on protease production, because growth was very low.

The effects of organic and inorganic nitrogen sources on protease production by *Bacillus* sp. have been reported in the literature. Many researchers have been reported that organic nitrogen sources was better suited to *Bacillus* sp. for growth and enzyme production than inorganic sources. It has been reported that peptone, casein, skim milk, yeast extract, favored maximum protease production by *Bacillus* sp (Puri et al. 2002; Sangeetha et al. 2008). Among the organic N sources, the effects of soybean flour, fish peptone, beef peptone and polypeptone were almost the same for *B. pumilus* c172-14 (Feng et al. 2001). The best nitrogen source for protease production was beef extract for *Bacillus* sp. K-30, while yeast extract and tryptone were comparable (Naidu and Devi 2005). Phadatare et al. (1993) reported the enhancement of protease production in *Conidiobolus coronatus* by organic nitrogen sources like tryptone, peptone and yeast extract. In the present study, addition of inorganic nitrogen sources in the production medium resulted in absence enzyme yield. It has also been reported by Fujiwara and Yamamoto (1987). On the other hand, some inorganic nitrogen sources gave better enzyme production. The best nitrogen source for protease production was $(\text{NH}_4)_2\text{SO}_4$ (Safey and Abdul-Raouf 2004). Among the various organic and inorganic nitrogen sources, the maximum enzyme activity was obtained with ammonium nitrate followed by ammonium chloride, ammonium citrate and potassium nitrate were used as nitrogen sources (Nascimento and Martins 2004).

The metal ions in media are an important factor that affects enzyme production due to act as inducers. The effects of some metal ions on protease activities were investigated. Firstly, Ca^{2+} and Mg^{2+} ions combined in culture medium (control) was tested along, and showed that enzyme production was very low. Hence, tested metal ions were used combine with each of Ca^{2+} and Mg^{2+} . Addition of Li^{2+} , Ba^{2+} , Mn^{2+} resulted in low production of protease. Cu^{2+} caused inhibition of production. Control (Ca^{2+} and Mg^{2+} combination) was better than tested metal ions (Fig. 2). It has been reported that Ca^{2+} and Mg^{2+} ions was effected on protease production (Feng et al. 2001; Kalaiarasi and Sunitha 2009). It suggested inhibitory effect of Cu^{2+} and Li^{2+} on proteases (Adinarayana et al. 2003).

Both our results and those of other reserachers have been shown that the protease production pathways of *Bacillus* strains were very different.

In the present study, the best carbon (fructose), nitrogen (skim milk) and metal ($\text{Ca}^{2+}+\text{Mg}^{2+}$) sources for protease production by *Bacillus* sp. N-40 were combined, and obtained a new medium. Production of protease in modified medium was realized. Maximum growth reached at 72 h, while maximum enzyme production (338 U/ml) was at 64 h. It was showed an increase of 51% in protease production when compared with the control (224 U/ml) (Fig. 3).




Figure 2. Protease production by *Bacillus* sp. N-40 in the presence of combined metal ions




Figure 3. Comparison of protease production by *Bacillus* sp. N-40 in control medium (Quadar et al., 2009) and newly modified medium

Some properties of partial purified protease enzyme

The enzyme was partially purified with a 2.1 fold increase in specific activity with a yield of 30%. Enzymatic properties, such as effect of pH, temperature and metal ions on the activity of the purified protease were performed. The optimum pH of enzyme was determined as pH 7.0 (Fig 4). 100% of enzyme activity was still detectable at pH 7.0 after 4 h at 37 °C. This might be the neutral protease. Because, the optima of most of the neutral protease have been reported in literature supported our finding (Basu et al. 2007; Merhep Dini et al. 2009). It was shown that the enzyme also gave high activity in the alkaline pH range 6.0-9.0 (Fig.4).

The optimum temperature of enzyme was 55°C (Fig. 5). Enzyme activity was stable with temperature within the range of 40°C to 70°C as seen in Fig.5. Enzyme was also still active at 80°C. In literature, optima temperature have been reported between 30-70°C for *Bacillus* sp. protease (Huang et al. 2006; Morya and Yaday 2010). A similar result, Ammar et al. (2003) reported that, the optimum temperature for thermostable purified protease enzyme was 55 °C. The thermal stability of protease was examined by pre incubating the enzyme at 55°C for 3h. The protease preserved 77% of its original activity.




Figure 4. Effect of pH on the protease activity




Figure 5. Effect of temperature on the protease activity

Metal ions are known to play a role as cofactors for enzyme activities, and often act as salt or ion bridges between two adjacent amino acid residues. In this study, various metal ions such as Li^{2+} , Zn^{2+} , Mg^{2+} , Ba^{2+} , Mn^{2+} , Ca^{2+} , Cu^{2+} at 5 mM concentration were tested for protease activation/inhibition effect, and the results are given in Table 2. Ca^{2+} and Mn^{2+} are found to have activating effect. Enzyme activity increased 28% , 26% in the presence of Mn^{2+} and Ca^{2+} as compared with control, respectively. Li^{2+} had low effect on protease activity. The activity was reduced in the presence of Ba^{2+} , Cu^{2+} , Mg^{2+} , Zn^{2+} . In the present work, strong activation/inhibition effect of metal ions on protease activity was not observed. Similarly, some investigators have also reported that protease activity was stimulated by Mn^{2+} and Ca^{2+} (Nascimento and Martins 2004; Jaswal and Kocher 2007). But, the activity was reduced in the presence of Mn^{2+} and Ca^{2+} (Basu et al. 2007; El-Hadj-Ali et al. 2007; Zambare et al. 2007; Merhep Dini et al. 2009) and increased by Zn^{2+} and Ba^{2+} (Jaswal and Kocher 2007). On the other hand, inhibition by Hg^{2+} , Cu^{2+} , Fe^{2+} , Mg^{2+} , Ba^{2+} , Zn^{2+} and Li^{2+} was reported by Bhatiya and Jadeja (2010).

Table 2. Effect of various metal ions (5mM) on *Bacillus* sp. N-40 protease activity. (Values are shown as means of triplicates.)

None	100
CaCl ₂	126
MgSO ₄	96
MnCl ₂	128
BaCl ₂	98
CuCl ₂	98
LiSO ₄	102
ZnSO ₄	70

All the experiments performed independently in triplicate and the results are presented as the mean of three.

This partial purified enzyme was electrophoresed on 10% (w/v) SDS-PAGE and a single band was observed. Using standard protein markers the size of the partially purified enzyme was found to be about 52 kDa (Fig. 6). A varieties of molecular weigth for proteases from other *Bacillus* species had been reported as 49 kDa *Bacillus* sp. HUTBS71 (Akel et al. 2009), 30,9 kDa *Bacillus* sp. HS08A (Huang et al. 2006), 75.0 kDa *Bacillus* sp. S17110 (Jung et al. 2007).




Figure 6. SDS-PAGE showing the molecular weight of protease enzyme produced by *Bacillus* sp. N-40 (52 kDa). A: S8445 SigmaMarker, B: Dialysate, C: Crude extract

CONCLUSIONS

A new strain of *Bacillus* was found to be a potential producer of protease enzyme. Studies on *Bacillus* sp. N-40 showed that nutritional factors include of sources of carbon, nitrogen and metal ions can inflence production of protease. Enzyme was partially purified, and characterized. This strain might be suitable for many industrial applications. Further experiments to enhanced enzyme production for commercialized process is needed.

REFERENCES

- Adinarayana K, Ellaiah P, and Prasad DS (2003). Purification and partial characterization of thermostable serine alkaline protease from a newly isolated *Bacillus subtilis* PE-11. AAPS Pharm. Sci. Tech., 4:1-9
- Akel H, Al-Quadan F, and Yousef TK (2009). Characterization of a Purified Thermostable Protease from Hyperthermophilic *Bacillus* Strain HUTBS71. Eur. J. Sci. Res., 31:280-288
- Ammar MS, Bayoumi RA, El-Kasaby AMH, and Soliman AM (2003). Purification and properties of thermostable protease by *B. brevis geltinoamylolyticus* using fish wastes (Fi.W.) and poultry wastes (Po.W) under solid state fermentation (S.S.F.) conditions 5th Int Sic Conf Al-Azhar Univ, Fac Sci Cairo, Egypt, pp 54
- Asokan S, and Jayanthi C (2010). Alkaline protease production by *Bacillus licheniformis* and *Bacillus coagulans*. J. Cell Tissue Res., 10: 2119-2123
- Basu BR, Banik AK, and Das M (2007). Production and characterization of extracellular protease of mutant *Aspergillus niger* AB100 grown on fish scale. World J. Microb. Biot. 24: 449-455
- Bhatiya R, and Jadeja GR (2010). Optimization of Environmental and Nutritional Factors for Alkaline Protease Production. EJEAFCh., 9: 594-599
- Boominadhan U, and Rajakumar R (2009). Optimization of Protease Enzyme Production Using *Bacillus* Sp. Isolated from Different Wastes. Bot. Res. Intl., 2:83-87
- Buchanan RE, and Gibbons NE (1974). Bergey's Manual of Determinative Bacteriology. 8thedn. The Williams and Wilkins Co., Baltimore
- El-Hadj-Ali N, Agrebi R, Ghorbel-Frikha B, Sellami-Kamoun A, S. Kanoun, and Nasri M (2007). Biochemical and molecular characterization of a detergent stable alkaline serine-protease from a newly isolated *Bacillus licheniformis* NH1. Enzyme Microb. Technol., 40:515-523
- El-Safey EM, and Abdul-Raouf UM (2004). Production, Purification and Characterization of Protease enzyme from *Bacillus subtilis*. International Conferences for Development and the Environment in the Arab World, Assiut University
- Feng YY, Yang WB, Ong SL, Hu JY, and Ng WJ (2001). Fermentation of starch for enhanced alkaline protease production by constructing an alkalophilic *Bacillus pumilus* Strain. Appl. Microbiol. Biotechnol., 57:153-160
- Fujiwara N, and Yamamoto K (1987). Production of alkaline protease in a low cost medium by alkalophilic *Bacillus* sp. and properties of the enzyme. J. Ferment. Technol., 65:345-348
- Fukushima Y, Itoh H, Fukasa T, and Motai H (1989). Continuous protease production in a carbon-limited chemostat culture by salt tolerant *Aspergillus oryzae*. Appl. Microbiol. Biotechnol., 30:604-608
- Gupta R, Beg QK, Khan S, and Chauhan B (2002). An overview on fermentation, downstream processing and properties of microbial alkaline proteases. Appl. Microbiol. Biotechnol., 60: 381-395
- Huang GR, Ying TJ, Huo P, and Jiang JX (2006). Purification and characterization of a protease from Thermophilic *Bacillus* strain HS08. Afr. J. Biotechnol., 5:2433-2438
- Jaswal RK, and Kocher GS (2007). Partial characterization of a crude alkaline protease from *Bacillus circulans* and its detergent compatibility. The Internet J. Microbiol., Vol. 4, No. 1.
- Johnvesly B, and Naik GR (2001). Studies on production of novel thermostable alkaline protease from thermophilic and alkaliphilic *Bacillus* sp. JB99 in a chemically defined medium. Process Biochem., 37:139-144
- Jung SC, Paik H, Kim MS, Baik KS, Lee W, Seong CN, and Choi SK (2007). InH_A-like protease secreted by *Bacillus* sp. S17110 inhabited in turban shell. J. Microbiol., 45:402-408
- Kalaiarasi K, and Sunitha PU (2009). Optimization of alkaline protease production from *Pseudomonas fluorescens* isolated from meat waste contaminated soil. Afr. J. Biotechnol., 8:7035-7041
- Keay L, and Wildi BS (1970). Proteinases of the genus *Bacillus*. I. Neutral proteinases. Biotechnol. Bioeng., 12:179-212
- Kim HK, Kim KH, Lee JK, Bae KS, Chang S, and Oh TK (1994). Purification and characterization of a thermostable protease from *Pseudomonas aeruginosa* NS-83. J. Microbiol. Biotechn., 4:113-118
- Laemmli UK (1970). Cleavages of structural proteins during the assembly of the head of Bacteriophage T4. Nature, 227: 680-685
- Lowry OH, Rosebrough NJ, Farr AL, and Randall RJ (1951). Protein measurement with the Folinphenol reagent. J. Biol. Chem., 193:265-275
- Mabrouk SS, Hashem AM, El-Shayeb NMA, Ismail AS, and Abdel-Fattah AF (1999). Optimization of alkaline protease productivity by *Bacillus licheniformis* ATCC 21415. Bioresour. Technol., 69: 155-159
- Mazdak C, Treton B, and Blanchin-Roland S (2000). Strong hybrid promoters and integrative expression/secretion vectors for quasi-constitutive expression of heterologous proteins in the yeast *Yarrowia lipolytica*. J. Mol. Microbiol. Biotechnol., 2:207-216
- McKellar RC, and Cholette H (1984). Synthesis of extracellular proteinase by *Pseudomonas fluorescens* under conditions of limiting carbon, nitrogen, and phosphate. Appl. Environ. Microbiol., 47:1224-1227
- Mehrato S, Pandey PK, Gaur R, and Darmwal NS (1999). The production of alkaline protease by a *Bacillus* species isolate. Bioresour. Technol., 67:201-203
- Merheb-Dini C, Cabral H, Leite RSR, Zanphorlin LM, Okamoto DN, Bonilla Rodriguez GO, Juliano L, Arantes EC, Gomes E, and Silva R (2009). Biochemical and functional characterization of a metalloprotease from the thermophilic fungus *Thermoascus aurantiacus*. J. Agric. Food. Chem., 57: 9210-9217
- Morya VK, and Yadav D (2010). Production and partial characterization of neutral protease by an indigenously isolated strain of *Aspergillus tubingensis* NIICC-08155. The Internet J. Microbiol., Vol. 8, No. 1
- Naidu KSB, and Devi KL (2005). Optimization of thermostable alkaline protease production from species of *Bacillus* using rice bran. Afr. J. Biotechnol., 4:724-726
- Nascimento WCA, and Martins MLL (2004). Production and properties of an extracellular protease from thermophilic *Bacillus* sp. Braz. J. Microbiol., 35:91-96
- Phadatar SU, Deshpande VV, and Srinivasan MC (1993). High activity alkaline protease from *Conidiobolus caronatus* (NCL 86.8.20): Enzyme production and compatibility with commercial detergents. Enzyme Microb. Tech., 15:72-76
- Puri S, Beg QK, and Gupta R (2002). Optimization of alkaline protease production from *Bacillus* sp. using response surface methodology. Curr. Microbiol., 44:286-290

- Quadar SA, Shireen E, Iqbal S, and Anwar A (2009). Optimization of protease production from newly isolated strain of *Bacillus* sp. PCSIR EA-3. *Indian J. Biotechnol.*, 8:286-290
- Rao MB, Tanksale AM, Ghatge MS, and Deshpande VV (1998). Molecular and Biotechnological Aspects of Microbial Proteases. *Microbiol. Mol. Biol. Rev.*, 62:597-635
- Sangeetha R, Geetha A, and Arulpandi I (2008). Optimization of protease and lipase production by *Bacillus pumilus* SG 2 isolated from an industrial effluent. *The Internet J. Microbiol.*, Vol. 5, No. 2
- Shafee N, Aris SN, Rahman RNZA, Basri M, and Salleh AB (2005). Optimization of environmental and nutritional conditions for the production of alkaline protease by a newly isolated bacterium *Bacillus cereus* strain 146. *J. Sci. Res.*, 1:1-8
- Zambare VP, Nilegaonkar SS, and Kanekar PP (2007). Production of an alkaline protease by *Bacillus cereus* MCM B-326 and its application as a dehairing agent. *World J. Microbiol. Biotechnol.*, 23:1569-1574