



Lie symmetry analysis, conservation laws and exact solutions of the seventh-order time fractional Sawada–Kotera–Ito equation



Emrullah Yaşar^a, Yakup Yıldırım^a, Chaudry Masood Khalique^{b,*}

^aDepartment of Mathematics, Faculty of Arts and Sciences, Uludag University, 16059 Bursa, Turkey

^bInternational Institute for Symmetry Analysis and Mathematical Modelling, Department of Mathematical Sciences, North-West University, Mafikeng Campus, Private Bag X 2046, Mmabatho 2735, South Africa

ARTICLE INFO

Article history:

Received 27 April 2016

Accepted 6 June 2016

Available online 11 June 2016

Keywords:

Fractional Sawada–Kotera–Ito equation

Lie symmetry

Riemann–Liouville fractional derivative

Conservation laws

Exact solutions

ABSTRACT

In this paper Lie symmetry analysis of the seventh-order time fractional Sawada–Kotera–Ito (FSKI) equation with Riemann–Liouville derivative is performed. Using the Lie point symmetries of FSKI equation, it is shown that it can be transformed into a nonlinear ordinary differential equation of fractional order with a new dependent variable. In the reduced equation the derivative is in Erdelyi–Kober sense. Furthermore, adapting the Ibragimov's nonlocal conservation method to time fractional partial differential equations, we obtain conservation laws of the underlying equation. In addition, we construct some exact travelling wave solutions for the FSKI equation using the sub-equation method.

© 2016 The Authors. Published by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

1. Introduction

Fractional partial differential equations (FPDEs) appear in various research and engineering applications such as physics, biology, rheology, viscoelasticity, control theory, signal processing, systems identification and electrochemistry [1–8]. Recently, they have attracted considerable interest and there has been a significant theoretical development in this area.

It is well-known that in order to describe nonlinear physical phenomena finding exact solutions of nonlinear fractional partial differential equations (NLFPEs) is the key instrument. A physical phenomenon may depend not only on the time instant but also on the time history, which can be successfully modelled using the theory of derivatives and integrals of fractional order [1–8].

Very recently, several powerful methods have been developed in the literature for finding exact solutions of NLFPEs. Some of the most important methods found in the literature include the exp function method, the fractional sub-equation method, the first integral method, the (G'/G) -expansion method, and the Lie symmetry method [9–28].

Lie symmetry analysis is one of the most general and effective methods for obtaining exact solutions of nonlinear partial differential equations (PDEs). In the last few decades, Lie's method has

been described in a number of excellent textbooks and has been applied to a number of physical and engineering models. See for example [29–33] and references therein.

However, application of the Lie symmetry analysis to FPDEs is quite new. We observe only a few studies in the literature. For instance, the authors of [15] considered the time fractional linear wave-diffusion equation and obtained a group of dilations. Using dilation symmetries invariant solutions were constructed. In [16], an attempt has been made to extend the Lie symmetry analysis to FPDEs (see also [17]). In addition, in [18], this method has been applied to time fractional generalized Burgers and Korteweg–de Vries equations. In [19], group-analysis of time fractional Harry–Dym equation with Riemann–Liouville derivative was performed and symmetry reductions and group-invariant solutions were obtained. The authors of [20] investigated the invariance properties of fractional Sharma–Tasso–Olver equation using the Lie group analysis method. In [21] an algorithm for the symbolic computation of Lie point symmetries for fractional differential equations (FDEs) was presented.

On the other hand, as stated in [30] conservation laws are very important tools in the study of differential equations from mathematical as well as physical point of view. If the underlying system has conservation laws then its integrability is quite possible [29,32]. Noether theorem [34] provides us with a systematic method for finding conservation laws of PDEs provided a Noether symmetry associated with a Lagrangian is known for Euler–Lagrange equations. However, there exist some approaches in the

* Corresponding author.

E-mail addresses: eyasar@uludag.edu.tr (E. Yaşar), yyildirim@windowslive.com (Y. Yıldırım), Masood.Khalique@nwu.ac.za (C.M. Khalique).

literature for obtaining the conservation laws of the PDEs, which do not have a Lagrangian [35–39].

The family of seventh-order Korteweg–de Vries (KdV) equations are given by [40]

$$u_t + au^3u_x + bu_x^3 + cuu_xu_{xx} + du^2u_{xxx} + eu_{2x}u_{3x} + fu_xu_{4x} + guu_{5x} + u_{7x} = 0, \tag{1}$$

where a, b, c, d, e, f and g are non-zero constants. In fact, the seventh-order KdV was introduced by Pomeau et al. [41] and its structural stability was discussed under a singular perturbation. In this paper, we study the seventh-order time fractional Sawada–Kotera–Ito (FSKI) equation

$$\frac{\partial^\alpha u}{\partial t^\alpha} + 252u^3u_x + 63u_x^3 + 378uu_xu_{xx} + 126u^2u_{xxx} + 63u_{2x}u_{3x} + 42u_xu_{4x} + 21uu_{5x} + u_{7x} = 0, \tag{2}$$

where $\alpha(0 < \alpha \leq 1)$ is a parameter describing the order of the fractional time derivative. When $\alpha = 1$, Eq. (2) becomes

$$\frac{\partial u}{\partial t} + 252u^3u_x + 63u_x^3 + 378uu_xu_{xx} + 126u^2u_{xxx} + 63u_{2x}u_{3x} + 42u_xu_{4x} + 21uu_{5x} + u_{7x} = 0, \tag{3}$$

which has been widely studied in the literature. For instance, the N -soliton solutions, bilinear form and Lax pair for Eq. (3) have been investigated in [42]. The bilinear and tanh–coth methods have been applied to Eq. (3) and soliton solutions were obtained in [43]. A decomposition was implemented for approximating its solutions [44]. In [45], using the Bell polynomial approach, Lax pair and infinite conservation laws were deduced for Eq. (3).

Our aim in the present work is to study symmetry reductions and conservation laws of the time FSKI equation (2) with the help of Lie symmetry analysis and Ibragimov’s nonlocal conservation method [38], respectively. In addition, we intend to obtain exact travelling wave solutions of the FSKI equation by the sub-equation method.

The paper is organized as follows. In Section 2, firstly some basic properties of Riemann–Liouville derivative are recalled. Then, Lie group method for FPDEs are presented. In Section 3, we apply the Lie group analysis method to the time FSKI equation (2) and obtain symmetry reductions. Then in Section 4, conservation laws of the time FSKI equation(2) are derived using the Ibragimov’s nonlocal conservation theorem. In Section 5, we construct exact travelling wave solutions of the time FSKI equation (2) via the sub-equation method. Lastly, concluding remarks are given in Section 6.

2. Preliminaries

We recall that the Riemann–Liouville derivative [4,5] of order α is defined by the following expression:

$$\partial_t^\alpha f(t) = \begin{cases} \frac{1}{\Gamma(n-\alpha)} \frac{\partial^n}{\partial t^n} \int_0^t \frac{f(\tau)}{(t-\tau)^{\alpha+1-n}} d\tau, & n-1 < \alpha < n, n \in \mathbb{N}, \\ \frac{\partial^n f}{\partial t^n}, & \alpha = n \in \mathbb{N}. \end{cases} \tag{4}$$

2.1. Description of Lie symmetry method for the time FPDEs

In this section, we present some notations and definitions that will be used in the sequel. For details see for example [16–20].

Consider a scalar time FPDE having the form [18,20]

$$E \equiv \frac{\partial^\alpha u}{\partial t^\alpha} - F(x, t, u, u_x, u_{xx}, u_{xxx}, u_{4x}, u_{5x}, u_{6x}, u_{7x}) = 0, \tag{5}$$

where $\alpha(0 < \alpha \leq 1)$ is a parameter. Consider a one-parameter Lie group of infinitesimal transformations given by

$$\bar{t} = t + \epsilon\tau(x, t, u) + O(\epsilon^2),$$

$$\bar{x} = x + \epsilon\xi(x, t, u) + O(\epsilon^2),$$

$$\bar{u} = u + \epsilon\eta(x, t, u) + O(\epsilon^2),$$

$$\frac{\partial^2 \bar{u}}{\partial \bar{t}^2} = \frac{\partial^2 u}{\partial t^2} + \epsilon\eta_x^0(x, t, u) + O(\epsilon^2),$$

$$\frac{\partial \bar{u}}{\partial \bar{x}} = \frac{\partial u}{\partial x} + \epsilon\eta^x(x, t, u) + O(\epsilon^2),$$

$$\frac{\partial^2 \bar{u}}{\partial \bar{x}^2} = \frac{\partial^2 u}{\partial x^2} + \epsilon\eta^{xx}(x, t, u) + O(\epsilon^2),$$

$$\frac{\partial^3 \bar{u}}{\partial \bar{x}^3} = \frac{\partial^3 u}{\partial x^3} + \epsilon\eta^{xxx}(x, t, u) + O(\epsilon^2),$$

$$\frac{\partial^4 \bar{u}}{\partial \bar{x}^4} = \frac{\partial^4 u}{\partial x^4} + \epsilon\eta^{xxxx}(x, t, u) + O(\epsilon^2),$$

$$\frac{\partial^5 \bar{u}}{\partial \bar{x}^5} = \frac{\partial^5 u}{\partial x^5} + \epsilon\eta^{xxxxx}(x, t, u) + O(\epsilon^2),$$

$$\frac{\partial^6 \bar{u}}{\partial \bar{x}^6} = \frac{\partial^6 u}{\partial x^6} + \epsilon\eta^{xxxxxx}(x, t, u) + O(\epsilon^2),$$

$$\frac{\partial^7 \bar{u}}{\partial \bar{x}^7} = \frac{\partial^7 u}{\partial x^7} + \epsilon\eta^{xxxxxxx}(x, t, u) + O(\epsilon^2), \tag{6}$$

where

$$\eta^x = D_x(\eta) - u_x D_x(\xi) - u_t D_x(\tau),$$

$$\eta^{xx} = D_x(\eta^x) - u_{xt} D_x(\tau) - u_{xx} D_x(\xi),$$

$$\eta^{xxx} = D_x(\eta^{xx}) - u_{xxt} D_x(\tau) - u_{xxx} D_x(\xi),$$

$$\eta^{xxxx} = D_x(\eta^{xxx}) - u_{xxxt} D_x(\tau) - u_{xxxx} D_x(\xi),$$

$$\eta^{xxxxx} = D_x(\eta^{xxxx}) - u_{xxxxt} D_x(\tau) - u_{xxxxx} D_x(\xi),$$

$$\eta^{xxxxxx} = D_x(\eta^{xxxxx}) - u_{xxxxxt} D_x(\tau) - u_{xxxxxx} D_x(\xi),$$

$$\eta^{xxxxxxx} = D_x(\eta^{xxxxxx}) - u_{xxxxxxt} D_x(\tau) - u_{xxxxxxx} D_x(\xi)$$

and D_x denotes the total differentiation operator defined by

$$D_x = \frac{\partial}{\partial x} + u_x \frac{\partial}{\partial u} + u_{xx} \frac{\partial}{\partial u_x} + \dots$$

Then the associated Lie algebra of symmetries is the set of vector fields of the form

$$X = \tau(x, t, u) \frac{\partial}{\partial t} + \xi(x, t, u) \frac{\partial}{\partial x} + \eta(x, t, u) \frac{\partial}{\partial u}. \tag{7}$$

The vector field (7) is a Lie point symmetry of (5) provided

$$\text{pr}^{(7)}X(E)|_{E=0} = 0. \tag{8}$$

Also, the invariance condition yields [20]

$$\tau(x, t, u)|_{t=0} = 0 \tag{9}$$

and the α th extended infinitesimal related to Riemann–Liouville fractional time derivative with (9) is given by [16,17]

$$\eta_x^0 = \frac{\partial^\alpha \eta}{\partial t^\alpha} + (\eta_u - \alpha D_t(\tau)) \frac{\partial^\alpha u}{\partial t^\alpha} - u \frac{\partial^\alpha \eta_u}{\partial t^\alpha} + \mu - \sum_{n=1}^{\infty} \binom{\alpha}{n} D_t^n(\xi) D_t^{\alpha-n}(u_x) + \sum_{n=1}^{\infty} \left[\binom{\alpha}{n} \frac{\partial^\alpha \eta_u}{\partial t^\alpha} - \binom{\alpha}{n+1} D_t^{n+1}(\tau) \right] D_t^{\alpha-n}(u), \tag{10}$$

where

$$\mu = \sum_{n=2}^{\infty} \sum_{m=2}^n \sum_{k=2}^m \sum_{r=0}^{k-1} \binom{\alpha}{n} \binom{n}{m} \binom{k}{r} \frac{1}{k!} \frac{t^{n-\alpha}}{\Gamma(n+1-\alpha)} [-u]^r \frac{\partial^m}{\partial t^m} \times [u^{k-r}] \frac{\partial^{n-m+k} \eta}{\partial t^{n-m} \partial u^k}. \tag{11}$$

It should be noted that we have $\mu = 0$ when the infinitesimal η is linear in u , because of the existence of the derivatives $\frac{\partial^k \eta}{\partial u^k}$, $k \geq 2$ in the above expression.

Definition 1. The function $u = \theta(x, t)$ is an invariant solution of Eq. (5) corresponding to the infinitesimal generator (7) if and only if

- (1) $u = \theta(x, t)$ satisfies Eq. (5).
- (2) $u = \theta(x, t)$ is an invariant surface of (6), namely, it fulfils the invariant surface condition

$$\tau(x, t, \theta) \theta_t + \xi(x, t, \theta) \theta_x = \eta(x, t, \theta).$$

3. Lie symmetries and reductions for the time FSKl equation

Let us assume that the time FSKl equation (2) is invariant under the one-parameter group of transformations (6) and so we have

$$\frac{\partial^\alpha \bar{u}}{\partial \bar{t}^\alpha} + 252\bar{u}^3 \bar{u}_x + 63\bar{u}_x^3 + 378\bar{u} \bar{u}_x \bar{u}_{xx} + 126\bar{u}^2 \bar{u}_{xxx} + 63\bar{u}_{2x} \bar{u}_{3x} + 42\bar{u}_x \bar{u}_{4x} + 21\bar{u} \bar{u}_{5x} + \bar{u}_{7x} = 0 \tag{12}$$

provided $u = u(x, t)$ satisfies (2). Using the point transformations (6) in (12) we obtain the invariant equation

$$\eta_x^0 + (756u^2 u_x + 378u_x u_{xx} + 252u u_{xxx} + 21u_{5x}) \eta + (252u^3 + 189u_x^2 + 378u u_{xx} + 42u_{4x}) \eta^x + (378u u_x + 63u_{3x}) \eta^{xx} + (126u^2 + 63u_{2x}) \eta^{xxx} + 42u_x \eta^{xxxx} + 21u \eta^{xxxxx} + \eta^{xxxxxx} = 0. \tag{13}$$

Substituting the values of $\eta_x^0, \eta^x, \eta^{xx}, \eta^{xxx}, \eta^{xxxx}, \eta^{xxxxx}$ and η^{xxxxxx} from (6) and (10) into (13) and equating various powers of the derivatives of u to zero, we obtain an overdetermined system of linear equations. These are (with the aid of [21])

$$\begin{aligned} \tau_x = \tau_u = \xi_t = \xi_u = \eta_{uu} = 0, \\ \binom{\alpha}{n} \partial_t^n (\eta_u) - \binom{\alpha}{n+1} D_t^{n+1} (\tau) = 0, \quad n = 1, 2, 3, \dots, \\ 7\xi'(x) - \alpha\tau'(t) = 0, \\ \partial_t^\alpha (\eta) - u \partial_t^\alpha (\eta_u) + 252u^3 \eta_x + 126u^2 \eta_{xxx} + 21u \eta_{xxxxx} + \eta_{xxxxxx} = 0. \end{aligned}$$

Now solving the above equations, we obtain

$$\xi = \alpha x c_1 + c_2, \quad \tau = 7t c_1, \quad \eta = -2\alpha u c_1,$$

where c_1 and c_2 are arbitrary constants. Hence the infinitesimal symmetry group of the time FSKl equation (2) is spanned by the two vector fields

$$X_1 = \frac{\partial}{\partial x}, \quad X_2 = 7t \frac{\partial}{\partial t} + \alpha x \frac{\partial}{\partial x} - 2\alpha u \frac{\partial}{\partial u}. \tag{14}$$

In what follows, we perform similarity reductions, present the reduced nonlinear fractional ordinary differential equations (ODEs) and classify the corresponding group-invariant solutions of the time FSKl equation (2) for the two Lie point symmetries (14).

Case 1. $X_1 = \partial/\partial x$. Integration of the invariant surface condition

$$\frac{dt}{0} = \frac{dx}{1} = \frac{du}{0}$$

gives the similarity variables t and u . Thus, we have the ansatz $u = f(t)$. Inserting this value of u into Eq. (2), we obtain the reduced fractional ODE

$$\partial_t^\alpha f(t) = 0.$$

Solving the above equation yields the group-invariant solution

$$u = a_1 t^{\alpha-1},$$

where a_1 is an arbitrary constant of integration.

Case 2. $X_2 = 7t \partial/\partial t + \alpha x \partial/\partial x - 2\alpha u \partial/\partial u$.

The similarity variables corresponding to the infinitesimal generator X_2 can be obtained by solving the associated characteristic equations given by

$$\frac{dt}{7t} = \frac{dx}{\alpha x} = \frac{du}{-2\alpha u}.$$

Solving the above equations, we obtain the two invariants

$$I_1 = ut^{2\alpha/7}, \quad I_2 = xt^{-\alpha/7}. \tag{15}$$

Thus, the symmetry X_2 gives the group-invariant solution

$$u = t^{-2\alpha/7} g(\xi), \quad \xi = xt^{-\alpha/7}, \tag{16}$$

where g is an arbitrary function of ξ . Using these invariants, Eq. (2) transforms to a special nonlinear ODE of fractional order. Thus, we have the following theorem corresponding to this case.

Theorem 1. The similarity transformation (16) reduces (2) to the following nonlinear ODE of fractional order:

$$\begin{aligned} \left(P_{\frac{7}{2}}^{1-\frac{9\alpha}{2}} g \right) (\xi) + 252g^3 g_\xi + 63g_\xi^3 + 378gg_\xi g_{\xi\xi} + 126g^2 g_{\xi\xi\xi} \\ + 63g_{2\xi} g_{3\xi} + 42g_\xi g_{4\xi} + 21gg_{5\xi} + g_{7\xi} = 0 \end{aligned} \tag{17}$$

with the Erdelyi–Kober fractional differential operator [4]

$$\left(P_\beta^{\tau, \alpha} g \right) := \prod_{j=0}^{n-1} \left(\tau + j - \frac{1}{\beta} \xi \frac{d}{d\xi} \right) \left(K_\beta^{\tau+\alpha, n-\alpha} g \right) (\xi), \tag{18}$$

$$n = \begin{cases} [\alpha] + 1, & \alpha \notin N, \\ \alpha, & \alpha \in N, \end{cases} \tag{19}$$

where

$$\left(K_\beta^{\tau, \alpha} g \right) (\xi) := \begin{cases} \frac{1}{\Gamma(\alpha)} \int_1^\infty (u-1)^{\alpha-1} u^{-(\tau+\alpha)} g(\xi u^{1/\beta}) du, & \alpha > 0, \\ g(\xi), & \alpha = 0, \end{cases} \tag{20}$$

is the Erdelyi–Kober fractional integral operator.

(See also [18,20]). Let $n-1 < \alpha < n$, $n = 1, 2, 3, \dots$. Based on the Riemann–Liouville fractional derivative for the similarity transformation (16), we have

$$\frac{\partial^\alpha u}{\partial t^\alpha} = \frac{\partial^n}{\partial t^n} \left[\frac{1}{\Gamma(n-\alpha)} \int_0^t (t-s)^{n-\alpha-1} s^{-\frac{2\alpha}{7}} g(xs^{-\frac{\alpha}{7}}) ds \right]. \tag{21}$$

Letting $v = t/s$, one can get $ds = -(t/v^2)dv$. Then Eq. (21) can be written as

$$\frac{\partial^\alpha u}{\partial t^\alpha} = \frac{\partial^n}{\partial t^n} \left[t^{n-\frac{9\alpha}{7}} \frac{1}{\Gamma(n-\alpha)} \int_1^\infty (v-1)^{n-\alpha-1} v^{-(n+1-\frac{9\alpha}{7})} g(\xi v^{\frac{\alpha}{7}}) dv \right]. \tag{22}$$

If one uses the definition of Erdelyi–Kober fractional integral operator (20), then Eq. (22) becomes

$$\frac{\partial^\alpha u}{\partial t^\alpha} = \frac{\partial^n}{\partial t^n} \left[t^{n-\frac{9\alpha}{7}} \left(K_{\frac{7}{2}}^{1-\frac{2\alpha}{7}, n-\alpha} g \right) (\xi) \right]. \tag{23}$$

We now intend to simplify the right hand side of (23). Taking into consideration $\xi = xt^{\frac{\alpha}{1-\alpha}}$, $\phi \in C^1(0, \infty)$, we can obtain

$$t \frac{\partial}{\partial t} \phi(\xi) = t x \left(-\frac{\alpha}{7}\right) t^{-\frac{\alpha}{7}-1} \phi'(\xi) = -\frac{\alpha}{7} \xi \frac{\partial}{\partial \xi} \phi(\xi). \tag{24}$$

Thus, we have

$$\begin{aligned} \frac{\partial^n}{\partial t^n} \left[t^{n-\frac{9\alpha}{7}} \left(K_{\frac{7}{2}}^{1-\frac{2\alpha}{7}, n-\alpha} g \right) (\xi) \right] &= \frac{\partial^{n-1}}{\partial t^{n-1}} \left[\frac{\partial}{\partial t} \left(t^{n-\frac{9\alpha}{7}} \left(K_{\frac{7}{2}}^{1-\frac{2\alpha}{7}, n-\alpha} g \right) (\xi) \right) \right] \\ &= \frac{\partial^{n-1}}{\partial t^{n-1}} \left[t^{n-\frac{9\alpha}{7}-1} \left(n - \frac{9\alpha}{7} - \frac{\alpha}{7} \xi \frac{\partial}{\partial \xi} \right) \left(K_{\frac{7}{2}}^{1-\frac{2\alpha}{7}, n-\alpha} g \right) (\xi) \right]. \end{aligned} \tag{25}$$

Repeating the same procedure $n - 1$ times, one can obtain

$$\begin{aligned} \frac{\partial^n}{\partial t^n} \left[t^{n-\frac{9\alpha}{7}} \left(K_{\frac{7}{2}}^{1-\frac{2\alpha}{7}, n-\alpha} g \right) (\xi) \right] &= \frac{\partial^{n-1}}{\partial t^{n-1}} \left[\frac{\partial}{\partial t} \left(t^{n-\frac{9\alpha}{7}} \left(K_{\frac{7}{2}}^{1-\frac{2\alpha}{7}, n-\alpha} g \right) (\xi) \right) \right] \\ &= \frac{\partial^{n-1}}{\partial t^{n-1}} \left[t^{n-\frac{9\alpha}{7}-1} \left(n - \frac{9\alpha}{7} - \frac{\alpha}{7} \xi \frac{\partial}{\partial \xi} \right) \left(K_{\frac{7}{2}}^{1-\frac{2\alpha}{7}, n-\alpha} g \right) (\xi) \right] \\ &\vdots \\ &= t^{-\frac{9\alpha}{7}} \prod_{j=0}^{n-1} \left(1 - \frac{9\alpha}{7} + j - \frac{\alpha}{7} \xi \frac{d}{d\xi} \right) \left(K_{\frac{7}{2}}^{1-\frac{2\alpha}{7}, n-\alpha} g \right) (\xi). \end{aligned} \tag{26}$$

Using the definition of the Erdélyi–Kober fractional differential operator (18) in the above, we get

$$\frac{\partial^n}{\partial t^n} \left[t^{n-\frac{9\alpha}{7}} \left(K_{\frac{7}{2}}^{1-\frac{2\alpha}{7}, n-\alpha} g \right) (\xi) \right] = t^{-\frac{9\alpha}{7}} \left(P_{\frac{7}{2}}^{1-\frac{9\alpha}{7}, \alpha} g \right) (\xi). \tag{27}$$

Now substituting (27) into (23), we obtain an expression for the time fractional derivative

$$\frac{\partial^\alpha u}{\partial t^\alpha} = t^{-\frac{9\alpha}{7}} \left(P_{\frac{7}{2}}^{1-\frac{9\alpha}{7}, \alpha} g \right) (\xi).$$

Thus, the time FSKL equation (2) can be reduced into a fractional order ODE

$$\begin{aligned} \left(P_{\frac{7}{2}}^{1-\frac{9\alpha}{7}, \alpha} g \right) (\xi) + 252g^3g_\xi + 63g_\xi^3 + 378gg_\xi g_{\xi\xi} + 126g^2g_{\xi\xi\xi} \\ + 63g_{2\xi}g_{3\xi} + 42g_\xi g_{4\xi} + 21gg_{5\xi} + g_{7\xi} = 0. \end{aligned} \tag{28}$$

This completes the proof of the theorem.

4. Conservation laws

We now construct the conservation laws of the FSKL equation (2). However, we first recall some basic definitions including the definitions of derivative and integral operators that we use in our work. The Riemann–Liouville left-sided time-fractional derivative

$${}_0D_t^\alpha u = D_t^n ({}_0I_t^{n-\alpha} u)$$

and the left-sided time-fractional integral of order $n - \alpha$, namely, ${}_0I_t^{n-\alpha}$ defined by

$$({}_0I_t^{n-\alpha} u)(x, t) = \frac{1}{\Gamma(n - \alpha)} \int_0^t \frac{u(\theta, x)}{(t - \theta)^{1-n+\alpha}} d\theta,$$

will be employed. Here $\Gamma(z)$ denotes the Gamma function, D_t is the operator of differentiation with respect to t and $n = [\alpha] + 1$ [27].

A conservation law for Eq. (2) is written as

$$D_t(C^t) + D_x(C^x) = 0,$$

which holds for all solutions $u(x, t)$ of the Eq. (2).

We now use Ibragimov method [38] for constructing the conservation laws of Eq. (2). It can easily be seen that the FSKL equation (2) has the formal Lagrangian

$$\begin{aligned} L = v(t, x) \left[\frac{\partial^2 u}{\partial t^2} + 252u^3u_x + 63u_x^3 + 378uu_xu_{xx} + 126u^2u_{xxx} \right. \\ \left. + 63u_{2x}u_{3x} + 42u_xu_{4x} + 21uu_{5x} + u_{7x} \right], \end{aligned}$$

where $v(t, x)$ is a new dependent variable. The Euler–Lagrange operator is [26,27]

$$\begin{aligned} \frac{\delta}{\delta u} = \frac{\partial}{\partial u} + (D_t^\alpha)^* \frac{\partial}{\partial D_t^\alpha u} - D_x \frac{\partial}{\partial u_x} + D_x^2 \frac{\partial}{\partial u_{xx}} - D_x^3 \frac{\partial}{\partial u_{xxx}} + D_x^4 \frac{\partial}{\partial u_{4x}} \\ - D_x^5 \frac{\partial}{\partial u_{5x}} - D_x^7 \frac{\partial}{\partial u_{7x}}, \end{aligned}$$

where $(D_t^\alpha)^*$ is the adjoint operator of (D_t^α) .

The adjoint equation to Eq. (2) is given by [38]

$$\frac{\delta L}{\delta u} = 0.$$

Also, we have [38]

$$\bar{X} + D_t(\tau)I + D_t(\xi)I = W \frac{\delta}{\delta u} + D_t N^t + D_x N^x,$$

where I is the identity operator, N^t and N^x are the Noether operators, and the prolonged vector field \bar{X} is defined by

$$\begin{aligned} \bar{X} = \tau \frac{\partial}{\partial t} + \xi \frac{\partial}{\partial x} + \eta \frac{\partial}{\partial u} + \eta_\alpha^0 \frac{\partial}{\partial D_t^\alpha u} + \eta^x \frac{\partial}{\partial u_x} + \eta^{xx} \frac{\partial}{\partial u_{xx}} + \eta^{xxx} \frac{\partial}{\partial u_{xxx}} \\ + \eta^{xxxx} \frac{\partial}{\partial u_{xxxx}} + \eta^{xxxxx} \frac{\partial}{\partial u_{xxxxx}} + \eta^{xxxxxx} \frac{\partial}{\partial u_{xxxxxx}} \end{aligned}$$

and the Lie characteristic function W is given by

$$W = \eta - \tau u_t - \xi u_x.$$

In the case when Riemann–Liouville time-fractional derivative is used in Eq. (2), the operator N^t is given by [26–28]

$$N^t = \tau I + \sum_{k=0}^{n-1} (-1)^k {}_0D_t^{\alpha-1-k} (W) D_t^k \frac{\partial}{\partial {}_0D_t^\alpha u} - (-1)^n J \left(W, D_t^n \frac{\partial}{\partial {}_0D_t^\alpha u} \right) \tag{29}$$

with J defined by

$$J(f, g) = \frac{1}{\Gamma(n - \alpha)} \int_0^t \int_t^T \frac{f(\tau, x)g(\mu, x)}{(\mu - \tau)^{\alpha+1-n}} d\mu d\tau. \tag{30}$$

For the seventh-order space derivatives, the operator N^x is given by

$$\begin{aligned} N^x = \xi I + W \left(\frac{\partial}{\partial u_x} - D_x \frac{\partial}{\partial u_{xx}} + D_x^2 \frac{\partial}{\partial u_{xxx}} - D_x^3 \frac{\partial}{\partial u_{4x}} + D_x^4 \frac{\partial}{\partial u_{5x}} - D_x^5 \frac{\partial}{\partial u_{6x}} + D_x^6 \frac{\partial}{\partial u_{7x}} \right) \\ + D_x(W) \left(\frac{\partial}{\partial u_{xx}} - D_x \frac{\partial}{\partial u_{xxx}} + D_x^2 \frac{\partial}{\partial u_{4x}} - D_x^3 \frac{\partial}{\partial u_{5x}} + D_x^4 \frac{\partial}{\partial u_{6x}} - D_x^5 \frac{\partial}{\partial u_{7x}} \right) \\ + D_x^2(W) \left(\frac{\partial}{\partial u_{xxx}} - D_x \frac{\partial}{\partial u_{4x}} + D_x^2 \frac{\partial}{\partial u_{5x}} - D_x^3 \frac{\partial}{\partial u_{6x}} + D_x^4 \frac{\partial}{\partial u_{7x}} \right) \\ + D_x^3(W) \left(\frac{\partial}{\partial u_{4x}} - D_x \frac{\partial}{\partial u_{5x}} + D_x^2 \frac{\partial}{\partial u_{6x}} - D_x^3 \frac{\partial}{\partial u_{7x}} \right) \\ + D_x^4(W) \left(\frac{\partial}{\partial u_{5x}} - D_x \frac{\partial}{\partial u_{6x}} + D_x^2 \frac{\partial}{\partial u_{7x}} \right) + D_x^5(W) \left(\frac{\partial}{\partial u_{6x}} - D_x \frac{\partial}{\partial u_{7x}} \right) \\ + D_x^6(W) \frac{\partial}{\partial u_{7x}}. \end{aligned} \tag{31}$$

The invariance condition for any given generator X of (2) and its solutions reads

$$(\bar{X}L + D_t(\tau)L + D_x(\xi)L)|_{(2)} = 0 \tag{32}$$

and consequently the conservation law of Eq. (2) can be written as

$$D_t(N^t L) + D_x(N^x L) = 0. \tag{33}$$

Now, we present the conservation laws of Eq. (2) using the above formalism. We consider two subcases corresponding to the order of α .

Case 1. When $\alpha \in (0, 1)$, with the help of (29) and (30), the components of the conserved vectors are

$$C_i^t = \tau L + (-1)^0 {}_0D_t^{\alpha-1}(W_i) D_t^0 \frac{\partial L}{\partial \{ {}_0D_t^\alpha u \}} - (-1)^1 J \left(W_i, D_t^1 \frac{\partial L}{\partial \{ {}_0D_t^\alpha u \}} \right) \\ = {}_0D_t^{\alpha-1}(W_i) v + J(W_i, v_t),$$

$$C_i^x = \xi L + W_i \left(\frac{\partial L}{\partial u_x} - D_x \frac{\partial L}{\partial u_{xx}} + D_x^2 \frac{\partial L}{\partial u_{xxx}} - D_x^3 \frac{\partial L}{\partial u_{4x}} + D_x^4 \frac{\partial L}{\partial u_{5x}} - D_x^5 \frac{\partial L}{\partial u_{6x}} + D_x^6 \frac{\partial L}{\partial u_{7x}} \right) \\ + D_x(W_i) \left(\frac{\partial L}{\partial u_{xx}} - D_x \frac{\partial L}{\partial u_{xxx}} + D_x^2 \frac{\partial L}{\partial u_{4x}} - D_x^3 \frac{\partial L}{\partial u_{5x}} + D_x^4 \frac{\partial L}{\partial u_{6x}} - D_x^5 \frac{\partial L}{\partial u_{7x}} \right) \\ + D_x^2(W_i) \left(\frac{\partial L}{\partial u_{xxx}} - D_x \frac{\partial L}{\partial u_{4x}} + D_x^2 \frac{\partial L}{\partial u_{5x}} - D_x^3 \frac{\partial L}{\partial u_{6x}} + D_x^4 \frac{\partial L}{\partial u_{7x}} \right) \\ + D_x^3(W_i) \left(\frac{\partial L}{\partial u_{4x}} - D_x \frac{\partial L}{\partial u_{5x}} + D_x^2 \frac{\partial L}{\partial u_{6x}} - D_x^3 \frac{\partial L}{\partial u_{7x}} \right) \\ + D_x^4(W_i) \left(\frac{\partial L}{\partial u_{5x}} - D_x \frac{\partial L}{\partial u_{6x}} + D_x^2 \frac{\partial L}{\partial u_{7x}} \right) + D_x^5(W_i) \left(\frac{\partial L}{\partial u_{6x}} - D_x \frac{\partial L}{\partial u_{7x}} \right) \\ + D_x^6(W_i) \frac{\partial L}{\partial u_{7x}}$$

$$= W_i \{ v(252u^3 + 189u_x^2 + 378uu_{xx} + 42u_{4x}) \\ - D_x(v(378uu_x + 63u_{3x})) + D_x^2(v(126u^2 + 63u_{2x})) - D_x^3(42vu_x) \\ + D_x^4(21vu) + D_x^5 v \} \\ + D_x(W_i) \{ v(378uu_x + 63u_{3x}) - D_x(v(126u^2 + 63u_{2x})) \\ + D_x^2(42vu_x) - D_x^3(21vu) - D_x^5 v \} \\ + D_x^2(W_i) \{ v(126u^2 + 63u_{2x}) - D_x(42vu_x) + D_x^2(21vu) + D_x^4 v \} \\ + D_x^3(W_i) \{ 42vu_x - D_x(21vu) - D_x^3 v \} + D_x^4(W_i) \{ 21vu + D_x^2 v \} \\ - v_x D_x^5(W_i) + v D_x^6(W_i),$$

where $i = 1, 2$ and the Lie characteristics functions W_i are given by $W_1 = -u_x, W_2 = -2\alpha u - 7tu_t - \alpha xu_x$.

Case 2. When $\alpha \in (1, 2)$, likewise as before the components of conserved vectors are given by

$$C_i^t = \tau L + (-1)^0 {}_0D_t^{\alpha-1}(W_i) D_t^0 \frac{\partial L}{\partial \{ {}_0D_t^\alpha u \}} - (-1)^1 J \left(W_i, D_t^1 \frac{\partial L}{\partial \{ {}_0D_t^\alpha u \}} \right) \\ + (-1)^1 {}_0D_t^{\alpha-2}(W_i) D_t^1 \frac{\partial L}{\partial \{ {}_0D_t^\alpha u \}} - (-1)^2 J \left(W_i, D_t^2 \frac{\partial L}{\partial \{ {}_0D_t^\alpha u \}} \right) \\ = v {}_0D_t^{\alpha-1}(W_i) + J(W_i, v_t) - v_t {}_0D_t^{\alpha-2}(W_i) - J(W_i, v_{tt}),$$

$$C_i^x = \xi L + W_i \left(\frac{\partial L}{\partial u_x} - D_x \frac{\partial L}{\partial u_{xx}} + D_x^2 \frac{\partial L}{\partial u_{xxx}} - D_x^3 \frac{\partial L}{\partial u_{4x}} + D_x^4 \frac{\partial L}{\partial u_{5x}} - D_x^5 \frac{\partial L}{\partial u_{6x}} + D_x^6 \frac{\partial L}{\partial u_{7x}} \right) \\ + D_x(W_i) \left(\frac{\partial L}{\partial u_{xx}} - D_x \frac{\partial L}{\partial u_{xxx}} + D_x^2 \frac{\partial L}{\partial u_{4x}} - D_x^3 \frac{\partial L}{\partial u_{5x}} + D_x^4 \frac{\partial L}{\partial u_{6x}} - D_x^5 \frac{\partial L}{\partial u_{7x}} \right) \\ + D_x^2(W_i) \left(\frac{\partial L}{\partial u_{xxx}} - D_x \frac{\partial L}{\partial u_{4x}} + D_x^2 \frac{\partial L}{\partial u_{5x}} - D_x^3 \frac{\partial L}{\partial u_{6x}} + D_x^4 \frac{\partial L}{\partial u_{7x}} \right) \\ + D_x^3(W_i) \left(\frac{\partial L}{\partial u_{4x}} - D_x \frac{\partial L}{\partial u_{5x}} + D_x^2 \frac{\partial L}{\partial u_{6x}} - D_x^3 \frac{\partial L}{\partial u_{7x}} \right) \\ + D_x^4(W_i) \left(\frac{\partial L}{\partial u_{5x}} - D_x \frac{\partial L}{\partial u_{6x}} + D_x^2 \frac{\partial L}{\partial u_{7x}} \right) + D_x^5(W_i) \left(\frac{\partial L}{\partial u_{6x}} - D_x \frac{\partial L}{\partial u_{7x}} \right) \\ + D_x^6(W_i) \frac{\partial L}{\partial u_{7x}}$$

$$= W_i \{ v(252u^3 + 189u_x^2 + 378uu_{xx} + 42u_{4x}) \\ - D_x(v(378uu_x + 63u_{3x})) \\ + D_x^2(v(126u^2 + 63u_{2x})) - D_x^3(42vu_x) + D_x^4(21vu) + D_x^6 v \} \\ + D_x(W_i) \{ v(378uu_x + 63u_{3x}) - D_x(v(126u^2 + 63u_{2x})) \\ + D_x^2(42vu_x) - D_x^3(21vu) - D_x^5 v \} \\ + D_x^2(W_i) \{ v(126u^2 + 63u_{2x}) - D_x(42vu_x) + D_x^2(21vu) + D_x^4 v \} \\ + D_x^3(W_i) \{ 42vu_x - D_x(21vu) - D_x^3 v \} + D_x^4(W_i) \{ 21vu + D_x^2 v \} \\ - v_x D_x^5(W_i) + v D_x^6(W_i),$$

where $i = 1, 2$ and functions W_i are given by

$$W_1 = -u_x, W_2 = -2\alpha u - 7tu_t - \alpha xu_x.$$

5. Exact travelling wave solutions of the time FSKI equation

In this section we construct exact travelling wave solutions of the time FSKI equation (2). For this aim, we consider the fractional derivative appears in the Eq. (2) in the sense of modified Riemann–Liouville derivative. The Jumarie’s modified Riemann–Liouville derivative [46] of order α is defined by the following expression

$$D_t^\alpha f(t) = \begin{cases} \frac{1}{\Gamma(1-\alpha)} \frac{d}{dt} \int_0^t (t-\zeta)^{-\alpha} (f(\zeta) - f(0)) d\zeta, & 0 < \alpha < 1, \\ [f^{(n)}(t)]^{(\alpha-n)}, & n \leq \alpha < n+1, n \geq 1. \end{cases} \tag{34}$$

where $f : R \rightarrow R, t \rightarrow f(t)$ denotes a continuous (but not necessarily first-order-differentiable) function and $\Gamma(\cdot)$ is the Gamma function defined by:

$$\Gamma(\alpha) = \lim_{n \rightarrow \infty} \frac{n! n^\alpha}{\alpha(\alpha+1)(\alpha+2) \dots (\alpha+n)} \tag{35}$$

Modified Riemann–Liouville derivative has the following important property:

$$D_t^\alpha t^\gamma = \frac{\Gamma(\gamma+1)}{\Gamma(\gamma+1-\alpha)} t^{\gamma-\alpha}, \quad \gamma > 0. \tag{36}$$

5.1. Description of the sub-equation method

In this subsection, we employ the sub-equation method, which was developed by Zhang and Zhang [12]. We recall this method here.

We consider the NLFPDEs of the type [12,20]

$$P(u, u_t, u_x, D_t^\alpha u, D_x^\alpha u, \dots) = 0, \quad 0 < \alpha \leq 1, \tag{37}$$

where u is an unknown function, F is a polynomial of u and its partial fractional derivatives, $D_x^\alpha u$ and $D_t^\alpha u$ are the modified Riemann–Liouville derivatives of u with respect to t and x , respectively. We present the main steps of the sub-equation method.

Step 1: By making use of the travelling wave transformation

$$u(x, t) = u(\zeta), \quad \zeta = x + ct, \tag{38}$$

where c is a nonzero constant to be determined later, we can rewrite (37) in the following nonlinear fractional ordinary differential equation (NFODE):

$$P(u, cu', u', c^\alpha D_\zeta^\alpha u, D_\zeta^\alpha u, \dots) = 0, \quad 0 < \alpha \leq 1. \tag{39}$$

Step 2: According to sub-equation method, we assume that the travelling wave solution can be expressed in the form

$$u(\zeta) = a_0 + \sum_{i=1}^n a_i (\phi(\zeta))^i, \tag{40}$$

where a_i ($i = 1, \dots, n$) are constants to be determined later, n is a positive integer which is determined by balancing the highest order derivatives and nonlinear terms in (37) and the function $\phi(\xi)$ satisfies the fractional Riccati equation

$$D_\xi^\alpha \phi(\xi) = \sigma + \phi^2(\xi), \tag{41}$$

where σ is a constant. Some special solutions of the fractional Riccati equation (41) are

$$\phi(\xi) = \begin{cases} -\sqrt{-\sigma} \tanh_x(\sqrt{-\sigma}\xi), & \sigma < 0, \\ -\sqrt{-\sigma} \coth_x(\sqrt{-\sigma}\xi), & \sigma < 0, \\ \sqrt{\sigma} \tan_x(\sqrt{\sigma}\xi), & \sigma > 0, \\ -\sqrt{\sigma} \cot_x(\sqrt{\sigma}\xi), & \sigma > 0, \\ -\frac{\Gamma(1+\alpha)}{\xi^\alpha + \omega}, & \omega = \text{const.}, \sigma = 0. \end{cases} \tag{42}$$

Step 3: Substituting (40) into (39) and setting the coefficients of ϕ^i to be zero, one obtains an over-determined nonlinear algebraic system in a_i ($i = 1, \dots, n$) and c .

Step 4: Then substituting these constants and the solutions of (41) into (40), we get the exact travelling solutions of the Eq. (37).

5.2. Application of the sub-equation method to the time FSKI equation

We now implement sub-equation method to Eq. (2). Firstly, we make use of the travelling wave transformation

$$u(x, t) = u(\xi), \quad \xi = x + ct,$$

where c is a constant and transform Eq. (2) into a nonlinear fractional ODE

$$c^\alpha D_\xi^\alpha u + 252u^3 u_\xi + 63u_\xi^3 + 378uu_\xi u_{\xi\xi} + 126u^2 u_{\xi\xi\xi} + 63u_{2\xi} u_{3\xi} + 42u_\xi u_{4\xi} + 21uu_{5\xi} + u_{7\xi} = 0. \tag{43}$$

We now suppose that Eq. (43) has the solution in the form

$$u(\xi) = a_0 + \sum_{i=1}^n a_i \phi(\xi)^i, \tag{44}$$

where a_i ($i = 1, \dots, n$) are constants to be determined and ϕ satisfies the fractional Riccati equation (41). Balancing the highest order derivative terms with nonlinear terms in Eq. (43), we get $n = 2$ and hence

$$u(\xi) = a_0 + a_1 \phi + a_2 \phi^2. \tag{45}$$

We then substitute Eq. (45) along with Eq. (41) into Eq. (43) then collect the coefficients of ϕ^i and equate them to zero. We obtain a set of algebraic equations in c, a_0, a_1 and a_2 . Solving these algebraic equations with the help of Maple, we get the following two cases:

Case 1:

$$a_1 = 0, \quad a_2 = -2, \quad c = (-252a_0^3 - 1008a_0^2\sigma - 1344a_0\sigma^2 - 608\sigma^3)^{\frac{1}{2}}.$$

Thus, from (42), we obtain five types of exact travelling wave solutions of Eq. (2), namely

$$\begin{aligned} u_1(x, t) &= a_0 - 2[-\sqrt{-\sigma} \tanh_x\{\sqrt{-\sigma}(x + ct)\}]^2, & \sigma < 0, \\ u_2(x, t) &= a_0 - 2[-\sqrt{-\sigma} \coth_x\{\sqrt{-\sigma}(x + ct)\}]^2, & \sigma < 0, \\ u_3(x, t) &= a_0 - 2[\sqrt{\sigma} \tan_x\{\sqrt{\sigma}(x + ct)\}]^2, & \sigma > 0, \\ u_4(x, t) &= a_0 - 2[-\sqrt{\sigma} \cot_x\{\sqrt{\sigma}(x + ct)\}]^2, & \sigma > 0, \\ u_5(x, t) &= a_0 - 2\left(-\frac{\Gamma(1+\alpha)}{(x + ct)^\alpha + \omega}\right)^2, & \omega = \text{const.}, \sigma = 0, \end{aligned}$$

where a_0 is an arbitrary constant.

Case 2:

$$a_0 = -8\sigma/3, \quad a_1 = 0, \quad a_2 = -4, \quad c = (-256\sigma^3/3)^{\frac{1}{2}}.$$

Likewise for this case, we also obtain five types of exact travelling solutions of Eq. (2) given by

$$\begin{aligned} u_1(x, t) &= \frac{-8\sigma}{3} - 4[-\sqrt{-\sigma} \tanh_x\{\sqrt{-\sigma}(x + ct)\}]^2, & \sigma < 0, \\ u_2(x, t) &= \frac{-8\sigma}{3} - 4[-\sqrt{-\sigma} \coth_x\{\sqrt{-\sigma}(x + ct)\}]^2, & \sigma < 0, \\ u_3(x, t) &= \frac{-8\sigma}{3} - 4[\sqrt{\sigma} \tan_x\{\sqrt{\sigma}(x + ct)\}]^2, & \sigma > 0, \\ u_4(x, t) &= \frac{-8\sigma}{3} - 4[-\sqrt{\sigma} \cot_x\{\sqrt{\sigma}(x + ct)\}]^2, & \sigma > 0, \\ u_5(x, t) &= \frac{-8\sigma}{3} - 4\left(-\frac{\Gamma(1+\alpha)}{(x + ct)^\alpha + \omega}\right)^2, & \omega = \text{const.}, \sigma = 0. \end{aligned}$$

6. Concluding remarks

In this work, we applied the Lie group analysis to time FSKI equation (2) based in the sense of Riemann–Liouville derivative. We obtained two– dimensional Lie symmetry algebra for (2). Using the nontrivial Lie point symmetry generator, we have shown that time FSKI equation can be transformed into a NFODE. Next, we constructed conservation laws for the time FSKI equation via Ibragimov’s nonlocal conservation theorem adapted to the time fractional partial differential equations. In addition, using the sub-equation method, we obtained hyperbolic, trigonometric and rational solutions to time FSKI equation (2). The exact solutions obtained here can be used as benchmarks against the numerical simulations. Furthermore, the exact solutions and conservation laws obtained in this paper might be very useful in various areas of applied mathematics in interpreting some physical phenomena.

References

- [1] Oldham KB, Spanier J. The fractional calculus. London: Academic Press; 1974.
- [2] Miller KS, Ross B. An introduction to the fractional calculus and fractional differential equations. New York: Wiley; 1993.
- [3] Samko S, Kilbas AA, Marichev O. Fractional integrals and derivatives: theory and applications. Yverdon: Gordon and Breach Science Publishers; 1993.
- [4] Kiryakova V. Generalised fractional calculus and applications. Pitman research notes in mathematics, vol. 301. London: Longman; 1994.
- [5] Podlubny I. Fractional differential equations. San Diego: Academic Press; 1999.
- [6] Shah K, Khalil H, Khan RA. Investigation of positive solution to a coupled system of impulsive boundary value problems for nonlinear fractional order differential equations. Chaos Solitons Fract 2015;77:240–6.
- [7] Biala TA, Jator SN. Block implicit Adams methods for fractional differential equations. Chaos Solitons Fract 2015;81:365–77.
- [8] Ahmad B, Ntouyas SK, Alsaedi A. On a coupled system of fractional differential equations with coupled nonlocal and integral boundary conditions. Chaos Solitons Fract 2016;83:234–41.
- [9] Bruzón MS, Garrido TM, de la Rosa R. Conservation laws and exact solutions of a Generalized Benjamin–Bona–Mahony–Burgers equation. Chaos Solitons Fract, to appear.
- [10] Recio E, Gandarias ML, Bruzón MS. Symmetries and conservation laws for a sixth–order Boussinesq equation. Chaos Solitons Fract, to appear.
- [11] Bekir A, Güner Ö, Çevikel AC. Fractional complex transform and exp-function methods for fractional differential equations. Abstr Appl Anal 2013. Article ID 426462, 8 p.
- [12] Zhang S, Zhang HQ. Fractional sub-equation method and its applications to nonlinear fractional PDEs. Phys Lett A 2011;375(7):1069–73.
- [13] Bekir A, Güner Ö, Ünsal Ö. The first integral method for exact solutions of nonlinear fractional differential equations. J Comput Nonlinear Dyn 2015;10. 021020–1.
- [14] Zheng B. (G/G)-expansion method for solving fractional partial differential equations in the theory of mathematical physics. Commun Theor Phys 2012;58:623–30.
- [15] Buckwar E, Luchko Y. Invariance of a partial differential equation of fractional order under the Lie group of scaling transformations. J Math Appl Math 1998;227:81–97.
- [16] Gazizov RK, Kasatkin AA, Lukashchuk SY. Continuous transformation groups of fractional differential equations. Vestnik, USATU 2007;9:125–35 (in Russian).

- [17] Gazizov RK, Kasatkin AA, Lukashchuk SY. Symmetry properties of fractional diffusion equations. *Phys Scr* 2009;T136:014016.
- [18] Sahadevan R, Bakkyaraj T. Invariant analysis of time fractional generalized Burgers and Korteweg–de Vries equations. *J Math Anal Appl* 2012;393:341–7.
- [19] Huang Q, Zhdanov R. Symmetries and exact solutions of the time fractional Harry–Dym equation with Riemann–Liouville derivative. *Physica A* 2014;409:110–8.
- [20] Wang GW, Xu TZ. Invariant analysis and exact solutions of nonlinear time fractional Sharma–Tasso–Olver equation by Lie group analysis. *Nonlinear Dyn* 2014;76:571–80.
- [21] Jefferson GF, Carminati J. FracSym: automated symbolic computation of Lie symmetries of fractional differential equations. *Comput Phys Commun* 2014;185:430–41.
- [22] Rui W, Zhang X. Lie symmetries and conservation laws for the time fractional Derrida–Lebowitz–Speer–Spohn equation. *Commun Nonlinear Sci Numer Simul* 2016;34:38–44.
- [23] Jafari H, Kadkhoda N, Baleanu D. Fractional Lie group method of the time-fractional Boussinesq equation. *Nonlinear Dyn* 2015;81(3):1569–74.
- [24] Bakkyaraj T, Sahadevan R. Invariant analysis of nonlinear fractional ordinary differential equations with Riemann–Liouville fractional derivative. *Nonlinear Dyn* 2015;80(1–2):447–55.
- [25] Hashemi MS. Group analysis and exact solutions of the time fractional Fokker–Planck equation. *Phys A: Stat Mech Appl* 2015;417:141–9.
- [26] Gazizov RK, Ibragimov NH, Lukashchuk SY. Nonlinear self-adjointness, conservation laws and exact solutions of time-fractional Kompaneets equations. *Commun Nonlinear Sci Numer Simul* 2014;23:153–63.
- [27] Lukashchuk SY. Conservation laws for time-fractional subdiffusion and diffusion-wave equations. *Nonlinear Dyn* 2015;80:791–802.
- [28] Wang G, Kara AH, Fakhri K. Symmetry analysis and conservation laws for the class of time-fractional nonlinear dispersive equation. *Nonlinear Dyn* 2015;82:281–7.
- [29] Bluman GW, Kumei S. Symmetries and differential equations. New York: Springer Verlag; 1989.
- [30] Olver PJ. Application of Lie groups to differential equations. New York: Springer-Verlag; 1993.
- [31] Ibragimov NH. Elementary Lie group analysis and ordinary differential equations. Chichester: John Wiley & Sons; 1999.
- [32] Adem AR, Khaliq CM. Symmetry reductions, exact solutions and conservation laws of a new coupled KdV system. *Commun Nonlinear Sci Numer Simul* 2012;17:3465–75.
- [33] Gandarias ML, Khaliq CM. Symmetries, solutions and conservation laws of a class of nonlinear dispersive wave equations. *Commun Nonlinear Sci Numer Simul* 2016;32:114–21.
- [34] Noether E. Invariante variationsprobleme. *Transp Theor Stat Phys* 1971;1(3):186–207. *Nachr. König. Gesell. Wissen., Göttingen. Math Phys Kl Heft* 1918;2:235–257. English translation.
- [35] Steudel H. Über die Zuordnung zwischen Invarianzeigenschaften und Erhaltungssätzen. *Z Naturforsch* 1962;17A:129–32.
- [36] Kara AH, Mahomed FM. Noether-type symmetries and conservation laws via partial Lagrangians. *Nonlinear Dyn* 2006;45:367–83.
- [37] Anco SC, Bluman GW. Direct construction method for conservation laws of partial differential equations. Part I: examples of conservation law classifications. *Eur J Appl Math* 2002;13:545–66.
- [38] Ibragimov NH. A new conservation theorem. *J Math Anal Appl* 2007;333:311–28.
- [39] Adem KR, Khaliq CM. Symmetry analysis and conservation laws of a generalized two-dimensional nonlinear KP–MEW equation. *Math Probl Eng* 2015;2015. <http://dx.doi.org/10.1155/2015/805763>. Article ID 805763, 7 p.
- [40] Göktaş Ü, Hereman W. Symbolic computation of conserved densities for systems of nonlinear evolution equations. *J Symbolic Comput* 1997;24:591–622.
- [41] Pomeau Y, Ramani A, Grammaticos B. Structural stability of the Korteweg–de Vries solitons under a singular perturbation. *Physica D* 1988;31:127–34.
- [42] Ito M. An extension of nonlinear evolution equations of the K–dV type to higher order. *J Phys Soc Jpn* 1980;49:771–8.
- [43] Wazwaz AM. The Hirota's direct method and the tanh-coth method for multiple-soliton solutions of the Sawada–Kotera–Ito seventh-order equation. *Appl Math Comput* 2008;199:133–8.
- [44] El-Sayed SM, Kaya D. An application of the ADM to seven-order Sawada–Kotera equations. *Appl Math Comput* 2004;157:93–101.
- [45] Shen YJ, Gao YT, Yu X, Meng GQ, Qin Y. Bell-polynomial approach applied to the seventh-order Sawada–Kotera–Ito equation. *Appl Math Comput* 2014;227:502–8.
- [46] Jumarie G. Modified Riemann–Liouville derivative and fractional Taylor series of nondifferentiable functions further results. *Comput Math Appl* 2006;51:1367–76.