



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Available online at
ScienceDirect
www.sciencedirect.com

Elsevier Masson France
EM|consulte
www.em-consulte.com



Original article

COVID-19: Initial experience of an international group of hand surgeons

Covid-19 : retour d'expérience d'un groupe international de chirurgiens de la main



F. Ducournau^a, M. Arianni^b, S. Awwad^c, E.-M. Baur^d, J.-Y. Beaulieu^e, M. Bouloudhine^f, M. Caloia^g, K. Chagar^h, Z. Chenⁱ, A.Y. Chin^j, E.C. Chow^k, T. Cobb^l, Y. David^m, P.J. Delgadoⁿ, M. Woon Man Fok^o, R. French^p, I. Golubev^q, J.R. Haugstvedt^r, S. Ichihara^s, R.A. Jorquera^t, S.C.J.J. Koo^u, J.Y. Lee^v, Y.K. Lee^w, Y.J. Lee^x, B. Liu^y, T. Kaleli^z, G.R. Mantovani^{aa}, C. Mathoulin^{ab}, J.C. Messina^{ac}, C. Muccioli^a, S. Nazerani^{ad}, C.Y. Ng^{ae}, M.C. Obdeijn^{af}, L. Van Overstraeten^{ag,ah}, T.O.H. Prasetyono^{ai}, M. Ross^{aj}, J.T. Shih^{ak}, N. Smith^{al}, F.A. Suarez R.^{am}, P.-T. Chan^{an}, H. Tiemdjo^{ao}, A. Wahegaonkar^{ap,aq}, M.C. Wells^{ar}, W.-Y. Wong^{as}, F. Wu^{at}, X.F. Yang^{au}, D. Yanni^{av}, J. Yao^{aw}, P.A. Liverneux^{a,*}

^a Icube CNRS 7357, department of hand surgery, hôpital de Hautepierre, university hospital of Strasbourg, FMFS, university of Strasbourg, 1, avenue Molière, 67200 Strasbourg cedex, France

^b Department of Surgery, RSUD Pasar Minggu Hospital, Jl. TB Simatupang No.1, RT.1/RW.5, Ragunan, Kec. Ps. Minggu, Kota Jakarta Selatan, Daerah Khusus Ibukota Jakarta 12550, Indonesia

^c National Guard Hospital Medina, Ad Dar, Medina, Saudi Arabia

^d Practice of Plastic and Hand Surgery, James-Loeb-Straße 13, 82418 Murnau am Staffelsee, Germany

^e Hôpitaux universitaires de Genève, département de chirurgie, rue Gabrielle-Perret-Gentil 4, 1205 Genève, Switzerland

^f Les Cliniques El Manar, 2092 rue Habib Echatti, Tunis, Tunisia

^g Department of Orthopaedic Surgery, Facultad de Ciencias Biomedicas, Universidad Austral, Hospital Universitario Austral, Pilar, Buenos Aires, Argentina

^h Pôle de chirurgie orthopédique, hôpital Militaire d'Instruction Mohamed V, Hay Riad, Rabat, Morocco

ⁱ Department of Hand Surgery, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, 328 Qi'an Ave, Xinzhou, Wuhan, Hubei, China

^j Department of Hand and Reconstructive Microsurgery, Singapore General Hospital, Academia Building, Outram Rd, Singapour 169608, Singapore

^k Department of Orthopaedics and Traumatology, United Christian Hospital, 130 Hip Wo St, Kwun Tong, Hong Kong

^l Orthopedic Specialists, P.C., 4622 Progress Drive Suite C, Davenport, IA 52807, USA

^m Hand surgery unit and microsurgery, Hospital Kaplan medical center, Derech Pasternak, Rehovot, Israel

ⁿ Hand Surgery and Microsurgery Department, University Hospital HM Montepincipe, CEU San Pablo University, Calle de Julián Romea 23, 28003 Madrid, Spain

^o Department of Orthopaedics and Traumatology 5/F, Professorial Block, Queen Mary Hospital, The University of Hong Kong, Queen Mary Hospital Main Block, Pok Fu Lam Rd, Pok Fu Lam, Hong Kong

^p The Specialist Referral Clinic, 555 W 12th Ave #121, Vancouver, BC V5Z 3X7, Canada

^q N. Priorov Research Institute of Trauma Surgery and Orthopaedics, 10 Moscow Ulitsa Priorova, Russia

^r Division of Handsurgery, Department of Orthopedics, Østfold Hospital, Per Gyntes vei 78, 1535 Moss, Norway

^s Department of Hand Surgery, Juntendo University Urayasu Hospital, 2 Chome-1-1 Tomioka, Urayasu, Chiba 279-0021, Japan

^t Department of Hand Surgery and Microsurgery, Clínica Indisa, Andrés Bello University, Av. Sta. María 1810, Santiago, Providencia, Región Metropolitana, Chile

^u Department of Orthopaedics and Traumatology, Alice Ho Miu Ling Nethersole Hospital, 11 Chuen On Road, Tai Po, NT, Hong Kong

^v Department of Orthopedic Surgery, The Catholic University of Korea, Eunpyeong St. Mary's Hospital, 1021 Tongil-ro, Eunpyeong-gu, 03312 Seoul, Korea

^w Department of Orthopedic Surgery, Research Institute of Clinical Medicine of Jeonbuk National University-Biomedical Research Institute of Jeonbuk National University Hospital, 567 Baekje-daero, Geumam 1(il)-dong, Deokjin-gu, Jeonju-si, Jeollabuk-do, Republic of Korea

^x Department of Orthopaedic Surgery, Kyungpook National University Hospital, 130 Dongdeok-ro, Jung-gu, Daegu 41944, Republic of Korea

^y Department of Hand Surgery, Beijing Ji Shui Tan Hospital, the 4th Clinical College of Peking University, 31 Xijiekou E Rd, Beijing Xicheng District, China

^z Uludag University, Faculty of Medicine, Department of Orthopaedics, Hand Surgery Clinic, Özlüce, Görükle Kampüsü, 16059 Nilüfer/Bursa, Turkey

^{aa} Department of Hand Surgery, Sao Paolo Hand center, Ben Portuguesa Hospital, R. Maestro Cardim 769, Bela Vista, São Paulo, SP, 01323-001, Brazil

^{ab} International Wrist Center, clinique Bizet, Institut de la main, 23, rue Georges-Bizet, 75116 Paris, France

^{ac} Gaetano Pini- CTO Orthopaedic Institute, First Orthopaedic Clinic University of Milan, Piazza Cardinale Andrea Ferrari 1, 20122 Milano MI, Italy

^{ad} Department of Hand and Reconstructive Microsurgery, Mehr General Hospital, W Zartosht St, District 6, Tehran, Tehran Province, IR, Iran

^{ae} Upper Limb Unit, Wrightington Hospital, Appley Bridge, Wigan, UK

^{af} Department of Plastic, Reconstructive and Hand surgery, Amsterdam University Medical Centers, AMC, Meibergdreef 9, 1105 AZ Amsterdam, The Netherlands

^{ag} Hôpital Erasme, ULB, route de Lennik 808, 1070 Bruxelles, Belgium

* Corresponding author.

E-mail address: Philippe.liverneux@chru-strasbourg.fr (P.A. Liverneux).

<https://doi.org/10.1016/j.hansur.2020.04.001>

2468-1229/© 2020 SFCM. Published by Elsevier Masson SAS. All rights reserved.

^{ah} Hand and foot surgery unit, Rue Pierre Caille 9, 7500 Tournai, Belgium

^{ai} Division of Plastic Surgery, Department of Surgery, Cipto Mangunkusumo Hospital/Faculty of Medicine Universitas Indonesia, Jl. Pangeran Diponegoro No.71, RW.5, Kenari, Kec. Senen, Kota Jakarta Pusat, Daerah Khusus Ibukota Jakarta 10430, Indonesia

^{aj} Brisbane Hand and Upper Limb Research Institute, Brisbane Private Hospital, 259 Wickham Terrace, Brisbane City QLD 4000, Australia

^{ak} Department of Orthopaedic Surgery, Armed Forces Taoyuan General Hospital, No. 168, Zhongxing Road, Longtan District, Taoyuan City, Taiwan

^{al} Southern Highlands Private Hospital, St Jude Specialist Centre, 21 St Jude St, Bowral NSW 2576, Australia

^{am} Universidad Militar Nueva Granada, Bogota, Colombia/Private clinic Centro de Cirugia Minimamente Invasiva CECIMIN, 76, Autopista Nte. #104, Bogotá, Colombia

^{an} Department of Orthopaedics and Traumatology, Tuen Mun Hospital, Block H, Tsing Chung Koon Rd, Tuen Mun, Hong Kong

^{ao} Centre de chirurgie de la main et des paralysies de Douala, Bonamoussadi, Douala, Cameroon

^{ap} Dept of Upper Limb, Hand and Microvascular Reconstructive Surgery, Brachial Plexus and Peripheral Nerve Surgery, Sancheti Institute for Orthopaedics and Rehabilitation, Jehangir Hospital, 32, Sasoon Road Opposite Railway Station, Central Excise Colony, Sangamvadi, Pune, Maharashtra 411001, India

^{aq} The Hand Surgery Clinics, 81/A/11 Giridarshan Society, Behind NEXA Showroom, Baner Road, Pune, Maharashtra 411007, India

^{ar} Orthopaedic hand surgeon, Mediclinic Panorama, Panorama, Rothschild Blvd, Panorama, Cape Town, 7500, South Africa

^{as} Department of Orthopaedics and Traumatology, the Chinese University of Hong Kong, Central Ave, Hong Kong

^{at} Dept of Orthopaedics, University Hospitals Birmingham, Bordesley Green East, Bordesley Green E, Birmingham B9 5SS, UK

^{au} Department of Hand Surgery, Union Hospital, Tongji Medical College, Huazhong University of Science and Technology, Jiang'an District, Wuhan, Hubei, China

^{av} KIMS Hospital, Newnham Ct Way, Weaving, Maidstone ME14 5FT, Kent, UK

^{aw} Robert A. Chase Hand and Upper Limb Center, Stanford University Medical Center, 450 Broadway, Redwood City, CA 94063, USA

ARTICLE INFO

Article history:

Received 27 March 2020

Received in revised form 1st April 2020

Accepted 5 April 2020

Available online 9 April 2020

Keywords:

COVID-19
Hand Surgery
Pandemic
Epidemic
Coronavirus

Mots clés :

Covid_19
Chirurgie de la main
Pandémie
Épidémie
Coronavirus

ABSTRACT

The emergence of the COVID-19 pandemic has severely affected medical treatment protocols throughout the world. While the pandemic does not affect hand surgeons at first glance, they have a role to play. The purpose of this study was to describe the different measures that have been put in place in response to the COVID-19 pandemic by hand surgeons throughout the world. The survey comprised 47 surgeons working in 34 countries who responded to an online questionnaire. We found that the protocols varied in terms of visitors, health professionals in the operating room, patient waiting areas, wards and emergency rooms. Based on these preliminary findings, an international consensus on hand surgery practices for the current viral pandemic, and future ones, needs to be built rapidly.

© 2020 SFCM. Published by Elsevier Masson SAS. All rights reserved.

R É S U M É

L'émergence de la pandémie à Covid-19 a bouleversé les pratiques médicales dans le monde. À première vue, cette pandémie ne concerne pas les chirurgiens de la main. Pourtant, ils ont un rôle à jouer. Le but de cette étude était de décrire les différentes pratiques mises en œuvre contre la pandémie à Covid-19 à partir du retour d'expérience de 47 chirurgiens de la main exerçant sur tous les continents. Le matériel comprenait 47 chirurgiens de la main exerçant dans 34 pays qui ont répondu à un questionnaire en ligne portant sur les protocoles Covid-19 mis en place. Les résultats ont montré que les pratiques étaient hétérogènes, tant pour les patients, les visiteurs et le personnel soignant au bloc opératoire, en salle de réunion, aux services d'hébergement des patients, au service d'accueil des urgences et en consultations externes. Il nous semble essentiel de définir un consensus international en chirurgie de la main pour lutter contre les pandémies virales actuelles et futures.

© 2020 SFCM. Publié par Elsevier Masson SAS. Tous droits réservés.

1. Introduction

The emergence of the COVID-19 pandemic has severely impacted medical practices throughout the world especially those of infectious disease specialists, emergency room personnel, anesthetists, intensivists, virologists, epidemiologists and hygienists. At first glance, the current pandemic does not affect hand surgeons; however, there is evidence that surgeons play a role in the continuation of emergency surgical care, the protection of medical staff and the management of resources [1].

A hand surgeon and an assistant from the Wuhan region were infected on 22nd January 2020 whilst operating on a COVID-19 patient. They self-isolated for 14 days. On the 3rd day, the surgeon showed signs of mild respiratory symptoms that got worse by the 9th day. The COVID-19 diagnosis was confirmed by CT scan and PCR (polymerase chain reaction) testing. Subsequently, the surgeon was admitted to a hospital where he eventually recovered. Review of this case showed that the protective measures in place against COVID-19 had not been followed correctly [2].

The purpose of this study was to describe the different measures that have been put in place in response to the COVID-19 pandemic by hand surgeons throughout the world.

2. Material and methods

The study comprised 47 hand surgeons (level 5 [3]) practicing in 34 different countries (Fig. 1). The survey consisted of an online questionnaire filled out between March 23 and 27, 2020. The questionnaire focused on the existing protocols in five key areas: operating theatres, surgeons' meeting rooms, wards, emergency rooms and external consultations. The results were summarized by describing the key features of the most common protocols used.

3. Results

In the operating theatre, 5/47 did not modify the indications for surgery, 3/47 totally stopped performing surgery, and 39/47 modified their practice. Within this last group, 28 operated on




Fig. 3. Question 2: If you only operate certain emergencies among those that you would have operated in normal times, which ones do you operate outside the operating room in a small emergency room/day surgery center under local anesthesia?




Fig. 4. Question 3: If you only operate certain emergencies, what do you do with those that you would have operated under normal circumstances and that you do not operate at all?




Fig. 5. Question 4: In the operating room.

emergencies only, whereas 5 continued scheduled surgical activities (Fig. 2). Within the group that continued to operate on emergency cases, 38 performed surgery in a small operating theatre on the patients they would normally have operated on in the main theatre (Fig. 3). Twenty-seven altered their indications for surgery and proposed conservative treatment (Fig. 4). And while 32 surgeons operated on COVID-19 positive patients, these operating theatres were specifically dedicated to COVID-19

patients. The majority wore specific personal protection equipment (PPE); however, the nature of this PPE varied (Fig. 5).

Most surgeons took measures to limit contact in meeting rooms (Fig. 6). On the wards, numerous measures were in place to limit transmission of the virus; these applied to patients, visitors and healthcare workers (Figs. 7 and 8). In the emergency rooms surveyed, most surgical residents assisted the emergency room staff in screening COVID-19 patients (Fig. 9). Consultation practices




Fig. 6. Question 5: Staff meetings.




Fig. 7. Question 6: Organization of hospitalization for patients.




Fig. 8. Question 7: Organization of hospitalization for staff.




Fig. 9. Question 8: In the emergency department.

varied: 5 saw patients as before while 42 modified their practice (Fig. 10). Amongst those who continued consultations, the majority took measures to screen patients and used personal protection measures against COVID-19 (Fig. 11).

4. Discussion

The management of a viral pandemic necessitates preventative measures to reduce intra-hospital transmission [4]. During surgery, drastic measures were put in place to facilitate the care of patients who are confirmed or suspected of being infected with COVID-19 and reduce the risk of intraoperative transmission to healthcare professionals and other patients. Some hospitals have set protocols related to negative pressure ventilation [5] and more

frequent air exchanges as well as limiting the numbers of circulating personnel in the operating theatre [6].

Our study has limitations. The first is that the severity and spread of COVID-19 were not recognized initially. The epidemic in China started in late December 2019 [7] and a pandemic was declared worldwide on 12 March 2020 [8]. Countries that were geographically distant from China lagged in putting into place any counter measures. The second limitation of the study lies in the fact that the severity of the outbreak was not the same in all countries [9]. The third limitation is that different countries took different counter measures, thus explaining the disparity of our survey results.

Our results show that the measures put in place by 47 hand surgeons were not homogeneous. As far as we know, the WHO has not issued any recommendations for the operating theatre related




Fig. 10. Question 9: Organization of outpatient consultations.




Fig. 11. Question 10: If you continue to consult.

to the COVID-19 pandemic. We feel that the following measures should be implemented strictly:

- all non-urgent surgeries should be deferred to a later date;
- in the operating theatre, all members of the surgical team should comply with exactly the same measures put in place by the infectious diseases department of their respective hospitals;

- all in-person meetings should be cancelled and be replaced by video conferencing;
- the wards should follow the same strict measures;
- office consultations should be deferred or cancelled [10].

If this proves impossible, then the same strict preventative measures should be followed [11].

Given the infection of a hand surgeon in Wuhan [2] and our study's findings, an international consensus is needed on the measures to be taken by hand surgeons during a viral pandemic.

Human and animal rights

The authors declare that the work described has not involved experimentation on humans or animals.

Informed consent and patient details

The authors declare that the work described does not involve patients or volunteers.

Disclosure of interest

The authors declare that they have no competing interest.

Funding

This work did not receive any grant from funding agencies in the public, commercial, or not-for-profit sectors.

Author contributions

All authors attest that they meet the current International Committee of Medical Journal Editors (ICMJE) criteria for Authorship.

References

- [1] Liang ZC, Wang W, Murphy D, Hui JHP. Novel Coronavirus and orthopaedic surgery: early experiences from Singapore. *J Bone Joint Surg Am* 2020;00. <http://dx.doi.org/10.2106/JBJS.20.00236> [e1(1-5)d], (in press).
- [2] Wang K, Wang Q, Chen Z. A case of hand surgeon infection of COVID-19 due to exposure to treating a suspect patient. *Chinese J Hand Surg* 2020;36. <http://dx.doi.org/10.3760/cma.j.cn311653-20200304-00096> (2020-03-08), Available from URL: <http://rs.yiigle.com/yufabiao/1184341.htm>.
- [3] Tang JB, Giddins G. Why and how to report surgeons' levels of expertise. *J Hand Surg Eur* 2016;41:365–6.
- [4] Centers for Disease Control and Prevention. Coronavirus Disease 2019 (COVID-19) Situation Summary. Available from URL: <https://www.cdc.gov/coronavirus/2019-ncov/summary.html>. (accessed March 2020).
- [5] Ti LK, Ang LS, Foong TW, Ng BSW. What we do when a COVID-19 patient needs an operation: operating room preparation and guidance. *Can J Anaesth* 2020. <http://dx.doi.org/10.1007/s12630-020-01617-4> [Epub ahead of print] PubMed PMID: 32144591.
- [6] Wong J, Goh QY, Tan Z, Lie SA, Tay YC, Ng SY, et al. Preparing for a COVID-19 pandemic: a review of operating room outbreak response measures in a large tertiary hospital in Singapore. *Can J Anaesth* 2020. <http://dx.doi.org/10.1007/s12630-020-01620-9> [Epub ahead of print] Review. PubMed PMID: 32162212.
- [7] World Health Organisation. Announces COVID-19 outbreak a pandemic. Available from URL: <http://www.euro.who.int/en/health-topics/health-emergencies/coronavirus-covid-19/news/news/2020/3/who-announces-covid-19-outbreak-a-pandemic>.
- [8] World Health Organisation 2020c. Statement on the second meeting of the International Health Regulations (2005) Emergency Committee regarding the outbreak of novel Coronavirus (2019-nCoV). Available from URL: [https://www.who.int/news-room/detail/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-\(2005\)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-\(2019-ncov\)](https://www.who.int/news-room/detail/30-01-2020-statement-on-the-second-meeting-of-the-international-health-regulations-(2005)-emergency-committee-regarding-the-outbreak-of-novel-coronavirus-(2019-ncov)). Accessed 30 January 2020.
- [9] World Health Organization. Coronavirus disease 2019 (COVID 19) Situation Report- 24 March 2020. Available from URL: https://www.who.int/docs/default-source/coronaviruse/situation-reports/20200324-sitrep-64-covid-19.pdf?sfvrsn=703b2c40_2. (accessed March 2020).
- [10] Hollander JE, Carr BG. Virtually perfect? Telemedicine for Covid-19. *N Engl J Med* 2020. <http://dx.doi.org/10.1056/NEJMp2003539> [Epub ahead of print] PubMed PMID: 32160451.
- [11] World Health Organisation. Infection prevention and control during health care when COVID-19 is suspected. Interim Guidance – 19 March 2020. Available from URL: [https://www.who.int/publications-detail/infection-prevention-and-control-during-health-care-when-novel-coronavirus-\(ncov\)-infection-is-suspected-20200125](https://www.who.int/publications-detail/infection-prevention-and-control-during-health-care-when-novel-coronavirus-(ncov)-infection-is-suspected-20200125).